

Algebraic cycles on a very special EPW sextic

ROBERT LATERVEER

ABSTRACT – Motivated by the Beauville–Voisin conjecture about Chow rings of powers of $K3$ surfaces, we consider a similar conjecture for Chow rings of powers of EPW sextics. We prove part of this conjecture for the very special EPW sextic studied by Donten–Bury et alii. We also prove some other results concerning the Chow groups of this very special EPW sextic, and of certain related hyperkähler fourfolds.

MATHEMATICS SUBJECT CLASSIFICATION (2010). Primary: 14C15, 14C25, 14C30. Secondary: 14J32, 14J35, 14J70, 14K99.

KEYWORDS. Algebraic cycles, Chow groups, motives, finite–dimensional motives, weak splitting property, weak Lefschetz conjecture for Chow groups, multiplicative Chow–Künneth decomposition, Bloch–Beilinson filtration, EPW sextics, hyperkähler varieties, $K3$ surfaces, abelian varieties, Calabi–Yau varieties.

1. Introduction

For a smooth projective variety X over \mathbb{C} , let $A^i(X) = CH^i(X)_{\mathbb{Q}}$ denote the Chow group of codimension i algebraic cycles modulo rational equivalence with \mathbb{Q} –coefficients. Intersection product defines a ring structure on $A^*(X) = \bigoplus_i A^i(X)$. In the case of $K3$ surfaces, this ring structure has an interesting property:

THEOREM 1.1 (Beauville–Voisin [8]). *Let S be a $K3$ surface. Let $D_i, D'_i \in A^1(S)$ be a finite number of divisors. Then*

$$\sum_i D_i \cdot D'_i = 0 \text{ in } A^2(S) \iff \sum_i D_i \cdot D'_i = 0 \text{ in } H^4(S, \mathbb{Q}).$$

Conjecturally, a similar property holds for self–products of $K3$ surfaces:

CONJECTURE 1.2 (Beauville–Voisin). *Let S be a $K3$ surface. For $r \geq 1$, let $D^*(S^r) \subset A^*(S^r)$ be the \mathbb{Q} –subalgebra generated by (the pullbacks of) divisors and the diagonal of S . The restriction of the cycle class map induces an injection*

$$D^i(S^r) \rightarrow H^{2i}(S^r, \mathbb{Q})$$

for all i and all r .

(cf. [53], [54], [56], [58] for extensions and partial results concerning conjecture 1.2.)

Beauville has asked which varieties have behaviour similar to theorem 1.1 and conjecture 1.2. This is the problem of determining which varieties verify the “weak splitting property” of [7]. We briefly state this problem here as follows:

PROBLEM 1.3 (Beauville [7]). *Find a nice class \mathcal{C} of varieties (containing K3 surfaces and abelian varieties), such that for any $X \in \mathcal{C}$, the Chow ring of X admits a multiplicative bigrading $A_{(*)}^*(X)$, with*

$$A^i(X) = \bigoplus_{j \geq 0} A_{(j)}^i(X) \text{ for all } i.$$

This bigrading should split the conjectural Bloch–Beilinson filtration, in particular

$$A_{hom}^i(X) = \bigoplus_{j \geq 1} A_{(j)}^i(X).$$

It has been conjectured that hyperkähler varieties are in \mathcal{C} [7, Introduction]. Also, not all Calabi–Yau varieties can be in \mathcal{C} [7, Example 1.7(b)]. An interesting novel approach of problem 1.3 (as well as a reinterpretation of theorem 1.1) is provided by the concept of *multiplicative Chow–Künneth decomposition* (cf. [43], [50], [44] and subsection 2.3 below).

In this note, we ask whether EPW sextics might be in \mathcal{C} . An EPW sextic is a special sextic $X \subset \mathbb{P}^5(\mathbb{C})$ constructed in [18]. EPW sextics are not smooth; however, a generic EPW sextic is a quotient $X = X_0/(\sigma_0)$, where X_0 is a smooth hyperkähler variety (called a double EPW sextic) and σ_0 is an anti-symplectic involution [35, Theorem 1.1], [36]. Quotient varieties behave like smooth varieties with respect to intersection theory with rational coefficients, so the following conjecture makes sense:

CONJECTURE 1.4. *Let X be an EPW sextic, and assume X is a quotient variety $X = X_0/G$ with X_0 smooth and $G \subset \text{Aut}(X_0)$ a finite group. Then $X \in \mathcal{C}$.*

There are two reasons why conjecture 1.4 is likely to be true: first, because an EPW sextic is a Calabi–Yau hypersurface (and these are probably in \mathcal{C}); secondly, because the hyperkähler variety X_0 should be in \mathcal{C} , and the involution σ_0 should behave nicely with respect to the bigrading on $A_{(*)}^*(X_0)$. Let us optimistically suppose conjecture 1.4 is true, and see what consequences this entails for the Chow ring of EPW sextics. We recall that Chow groups are expected to satisfy a weak Lefschetz property, according to a long-standing conjecture:

CONJECTURE 1.5 (Hartshorne [24]). *Let $X \subset \mathbb{P}^{n+1}(\mathbb{C})$ be a smooth hypersurface of dimension $n \geq 4$. Then the cycle class map*

$$A^2(X) \rightarrow H^4(X, \mathbb{Q})$$

is injective.

Conjecture 1.5 is notoriously open for all hypersurfaces of degree $d \geq n + 2$. Since quotient varieties behave in many ways like smooth varieties, it seems reasonable to expect that conjecture 1.5 extends to hypersurfaces that are quotient varieties. This would imply that an EPW sextic X as in conjecture 1.4 has $A_{hom}^2(X) = 0$. That is, conjecturally we have that

$$A^i(X) = A_{(0)}^i(X) \text{ for all } i \leq 2.$$

For any $r \geq 1$, let us now define

$$E^*(X^r) \subset A^*(X^r)$$

as the \mathbb{Q} -subalgebra generated by (pullbacks of) elements of $A^1(X)$ and $A^2(X)$ and the class of the diagonal of X . The above remarks imply a conjectural inclusion

$$E^*(X^r) \subset A_{(0)}^*(X^r) = A^*(X^r)/A_{hom}^*(X^r).$$

We thus arrive at the following concrete, falsifiable conjecture:

CONJECTURE 1.6. *Let X be an EPW sextic as in conjecture 1.4. Then restriction of the cycle class map*

$$E^i(X^r) \rightarrow H^{2i}(X^r, \mathbb{Q})$$

is injective for all i and all r .

Conjecture 1.6 is the analogon of conjecture 1.2 for EPW sextics; the role of divisors on the $K3$ surface is played by (the hyperplane section and) codimension 2 cycles on the sextic. The main result in this note provides some evidence for conjecture 1.6: we can prove it is true for 0–cycles and 1–cycles on one very special EPW sextic:

THEOREM (=theorem 4.7). *Let X be the very special EPW sextic of [16]. Let $r \in \mathbb{N}$. The restriction of the cycle class map*

$$E^i(X^r) \rightarrow H^{2i}(X^r, \mathbb{Q})$$

is injective for $i \geq 4r - 1$.

The very special EPW sextic of [16] (cf. section 2.7 below for a definition) is not smooth, but it is a “Calabi–Yau variety with quotient singularities”. The very special EPW sextic X is very symmetric; it is also remarkable for providing the only example known so far of a complete family of 20 pairwise incident planes in $\mathbb{P}^5(\mathbb{C})$ [16]. As resumed in theorem 2.28 below, the very special EPW sextic X is related to hyperkähler varieties in two different ways: (a) X is rationally dominated via a degree 2 map by the Hilbert scheme $S^{[2]}$ where S is a $K3$ surface of Picard number 20; (b) X admits a double cover that is the quotient of an abelian variety by a finite group of group automorphisms, and this quotient admits a hyperkähler resolution X_0 .

To prove theorem 4.7, we first prove (proposition 3.3) that the very special EPW sextic X has a multiplicative Chow–Künneth decomposition, in the sense of Shen–Vial [43], and so the Chow ring of X has a bigrading. Next, we establish (proposition 3.8) that

$$(1) \quad A^2(X) = A_{(0)}^2(X).$$

Both these facts are proven using description (b), via the theory of *symmetrically distinguished cycles* [37].

Note that equality (1) might be considered as evidence for conjecture 1.5 for X . In order to prove conjecture 1.5 for the very special EPW sextic X , it remains to prove that

$$A_{(0)}^2(X) \cap A_{hom}^2(X) \stackrel{??}{=} 0.$$

Likewise, in order to prove the full conjecture 1.6 for the very special EPW sextic X , it remains to prove that

$$A_{(0)}^i(X^r) \cap A_{hom}^i(X^r) \stackrel{??}{=} 0 \quad \text{for all } i, r.$$

We are not able to prove these equalities outside of the range $i \geq 4r - 1$; this is related to some of the open cases of Beauville’s conjecture on Chow rings of abelian varieties (remarks 4.4 and 4.8).

On the positive side, we establish a precise relation between the Chow ring of the very special EPW sextic X and the Chow ring of the hyperkähler fourfold X_0 mentioned in description (b) (theorem 4.9). This relation provides an alternative description of the splitting of the Chow ring of X_0 coming from a multiplicative Chow–Künneth decomposition (corollary 4.10). In proving this relation, we exploit description (a); a key ingredient in the proof is a strong version of the generalized Hodge conjecture for X and X_0 (proposition 3.1), which crucially relies on the fact that the $K3$ surface S has maximal Picard number.

We also obtain some results concerning Bloch’s conjecture (subsection 5.1), as well as a conjecture of Voisin (subsection 5.2), for the very special EPW sextic. The application to Bloch’s conjecture relies on description (b) (via the theory of symmetrically distinguished cycles), but also on description (a) (via the surjectivity result proposition 3.12).

We end this introduction with a challenge: can one prove theorem 4.7 for other (not very special) EPW sextics ?

CONVENTIONS. *In this note, the word variety will refer to a reduced irreducible scheme of finite type over \mathbb{C} . A subvariety is a (possibly reducible) reduced subscheme which is equidimensional.*

All Chow groups will be with rational coefficients: *we denote by $A_j X$ the Chow group of j –dimensional cycles on X with \mathbb{Q} –coefficients; for X smooth of dimension n the notations $A_j X$ and $A^{n-j} X$ will be used interchangeably.*

The notations $A_{\text{hom}}^j(X)$, $A_{\text{num}}^j(X)$, $A_{AJ}^j(X)$ will be used to indicate the subgroups of homologically trivial, resp. numerically trivial, resp. Abel–Jacobi trivial cycles. The contravariant category of Chow motives (i.e., pure motives with respect to rational equivalence as in [42], [34]) will be denoted \mathcal{M}_{rat} .

We will write $H^j(X)$ and $H_j(X)$ to indicate singular cohomology $H^j(X, \mathbb{Q})$, resp. Borel–Moore homology $H_j(X, \mathbb{Q})$.

2. Preliminary material

2.1 – Quotient varieties

DEFINITION 2.1. A projective quotient variety is a variety

$$X = Y/G,$$

where Y is a smooth projective variety and $G \subset \text{Aut}(Y)$ is a finite group.

PROPOSITION 2.2 (Fulton [22]). Let X be a projective quotient variety of dimension n . Let $A^*(X)$ denote the operational Chow cohomology ring. The natural map

$$A^i(X) \rightarrow A_{n-i}(X)$$

is an isomorphism for all i .

PROOF. This is [22, Example 17.4.10]. □

REMARK 2.3. It follows from proposition 2.2 that the formalism of correspondences goes through unchanged for projective quotient varieties (this is also noted in [22, Example 16.1.13]). We can thus consider motives $(X, p, 0) \in \mathcal{M}_{\text{rat}}$, where X is a projective quotient variety and $p \in A^n(X \times X)$ is a projector. For a projective quotient variety $X = Y/G$, one readily proves (using Manin’s identity principle) that there is an isomorphism

$$h(X) \cong h(Y)^G := (Y, \Delta_Y^G, 0) \text{ in } \mathcal{M}_{\text{rat}},$$

where Δ_Y^G denotes the idempotent $\frac{1}{|G|} \sum_{g \in G} \Gamma_g$.

2.2 – Finite–dimensionality

We refer to [32], [4], [34], [26], [30] for basics on the notion of finite–dimensional motive. An essential property of varieties with finite–dimensional motive is embodied by the nilpotence theorem:

THEOREM 2.4 (Kimura [32]). Let X be a smooth projective variety of dimension n with finite–dimensional motive. Let $\Gamma \in A^n(X \times X)$ be a correspondence which is numerically trivial. Then there is $N \in \mathbb{N}$ such that

$$\Gamma^{\circ N} = 0 \quad \in A^n(X \times X).$$

Actually, the nilpotence property (for all powers of X) could serve as an alternative definition of finite–dimensional motive, as shown by a result of Jannsen [30, Corollary 3.9]. Conjecturally, all smooth projective varieties have finite–dimensional motive [32]. We are still far from knowing this, but at least there are quite a few non–trivial examples:

REMARK 2.5. The following varieties have finite–dimensional motive: abelian varieties, varieties dominated by products of curves [32], $K3$ surfaces with Picard number 19 or 20 [38], surfaces not of general type with $p_g = 0$ [23, Theorem 2.11], certain surfaces of general type with $p_g = 0$ [23], [40], [55], Hilbert schemes of surfaces known to have finite–dimensional motive [13], generalized Kummer varieties [57, Remark 2.9(ii)], [21], threefolds with nef tangent bundle [27], [47, Example 3.16], fourfolds with nef tangent

bundle [28], log-homogeneous varieties in the sense of [12] (this follows from [28, Theorem 4.4]), certain threefolds of general type [49, Section 8], varieties of dimension ≤ 3 rationally dominated by products of curves [47, Example 3.15], varieties X with $A_{AJ}^i(X) = 0$ for all i [46, Theorem 4], products of varieties with finite-dimensional motive [32].

REMARK 2.6. It is an embarrassing fact that up till now, all examples of finite-dimensional motives happen to lie in the tensor subcategory generated by Chow motives of curves, i.e. they are “motives of abelian type” in the sense of [47]. On the other hand, there exist many motives that lie outside this subcategory, e.g. the motive of a very general quintic hypersurface in \mathbb{P}^3 [14, 7.6].

The notion of finite-dimensionality is easily extended to quotient varieties:

DEFINITION 2.7. Let $X = Y/G$ be a projective quotient variety. We say that X has finite-dimensional motive if the motive

$$h(Y)^G := (Y, \Delta_Y^G, 0) \in \mathcal{M}_{\text{rat}}$$

is finite-dimensional. (Here, Δ_Y^G denotes the idempotent $\frac{1}{|G|} \sum_{g \in G} \Gamma_g \in A^n(Y \times Y)$.)

Clearly, if Y has finite-dimensional motive then also $X = Y/G$ has finite-dimensional motive. The nilpotence theorem extends to this set-up:

PROPOSITION 2.8. *Let $X = Y/G$ be a projective quotient variety of dimension n , and assume X has finite-dimensional motive. Let $\Gamma \in A_{\text{num}}^n(X \times X)$. Then there is $N \in \mathbb{N}$ such that*

$$\Gamma^{\circ N} = 0 \in A^n(X \times X).$$

PROOF. Let $p: Y \rightarrow X$ denote the quotient morphism. We associate to Γ a correspondence $\Gamma_Y \in A^n(Y \times Y)$ defined as

$$\Gamma_Y := {}^t\Gamma_p \circ \Gamma \circ \Gamma_p \in A^n(Y \times Y).$$

By Lieberman’s lemma [47, Lemma 3.3], there is equality

$$\Gamma_Y = (p \times p)^* \Gamma \text{ in } A^n(Y \times Y),$$

and so Γ_Y is $G \times G$ -invariant:

$$\Delta_Y^G \circ \Gamma_Y \circ \Delta_Y^G = \Gamma_Y \text{ in } A^n(Y \times Y).$$

This implies that

$$\Gamma_Y \in \Delta_Y^G \circ A^n(Y \times Y) \circ \Delta_Y^G,$$

and so

$$\Gamma_Y \in \text{End}_{\mathcal{M}_{\text{rat}}}(h(Y)^G).$$

Since clearly Γ_Y is numerically trivial, and $h(Y)^G$ is finite-dimensional (by assumption), there exists $N \in \mathbb{N}$ such that

$$(\Gamma_Y)^{\circ N} = {}^t\Gamma_p \circ \Gamma \circ \Gamma_p \circ {}^t\Gamma_p \circ \dots \circ \Gamma_p = 0 \text{ in } A^n(Y \times Y).$$

Using the relation $\Gamma_p \circ {}^t\Gamma_p = d\Delta_X$, this boils down to

$$d^{N-1} {}^t\Gamma_p \circ \Gamma^{\circ N} \circ \Gamma_p = 0 \text{ in } A^n(Y \times Y).$$

From this, we deduce that also

$$\Gamma^{\circ N} = \frac{1}{d^{N+1}} \Gamma_p \circ \left(d^{N-1} {}^t\Gamma_p \circ \Gamma^{\circ N} \circ \Gamma_p \right) \circ {}^t\Gamma_p = 0 \text{ in } A^n(X \times X).$$

□

2.3 – MCK decomposition

DEFINITION 2.9 (Murre [33]). Let X be a projective quotient variety of dimension n . We say that X has a *CK decomposition* if there exists a decomposition of the diagonal

$$\Delta_X = \Pi_0 + \Pi_1 + \cdots + \Pi_{2n} \quad \text{in } A^n(X \times X),$$

such that the Π_i are mutually orthogonal idempotents and $(\Pi_i)_* H^*(X) = H^i(X)$.

REMARK 2.10. The existence of a CK decomposition for any smooth projective variety is part of Murre’s conjectures [33], [29]. If a quotient variety X has finite-dimensional motive, and the Künneth components are algebraic, then X has a CK decomposition (this can be proven just as [29], where this is stated for smooth X).

DEFINITION 2.11 (Shen–Vial [43]). Let X be a projective quotient variety of dimension n . Let $\Delta_{sm}^X \in A^{2n}(X \times X \times X)$ be the class of the small diagonal

$$\Delta_{sm}^X := \{(x, x, x) \mid x \in X\} \subset X \times X \times X.$$

An MCK decomposition of X is a CK decomposition $\{\Pi_i\}$ of X that is *multiplicative*, i.e. it satisfies

$$\Pi_k \circ \Delta_{sm}^X \circ (\Pi_i \times \Pi_j) = 0 \quad \text{in } A^{2n}(X \times X \times X) \quad \text{for all } i + j \neq k.$$

(NB: the acronym “MCK” is shorthand for “multiplicative Chow–Künneth”.)

REMARK 2.12. The small diagonal (seen as a correspondence from $X \times X$ to X) induces the *multiplication morphism*

$$\Delta_{sm}^X: h(X) \otimes h(X) \rightarrow h(X) \quad \text{in } \mathcal{M}_{\text{rat}}.$$

Suppose X has a CK decomposition

$$h(X) = \bigoplus_{i=0}^{2n} h^i(X) \quad \text{in } \mathcal{M}_{\text{rat}}.$$

By definition, this decomposition is multiplicative if for any i, j the composition

$$h^i(X) \otimes h^j(X) \rightarrow h(X) \otimes h(X) \xrightarrow{\Delta_{sm}^X} h(X) \quad \text{in } \mathcal{M}_{\text{rat}}$$

factors through $h^{i+j}(X)$.

The property of having an MCK decomposition is severely restrictive, and is closely related to Beauville’s “weak splitting property” [7]. For more ample discussion, and examples of varieties with an MCK decomposition, we refer to [43, Section 8] and also [50], [44], [21].

LEMMA 2.13. *Let X, X' be birational hyperkähler varieties. Then X has an MCK decomposition if and only if X' has one.*

PROOF. This is noted in [50, Introduction]; the idea is that Rieß’s result [41] implies that X and X' have isomorphic Chow motives and the isomorphism is compatible with the multiplicative structure.

More precisely: let $\phi: X \dashrightarrow X'$ be a birational map between hyperkähler varieties of dimension n . According to [41] there exists a correspondence $\gamma \in A^n(X \times X')$ inducing a ring isomorphism $A^*(X) \cong A^*(X')$.

Suppose that $\{\Pi_i^X\}$ is an MCK decomposition for X . Let $\Delta_{sm}^X, \Delta_{sm}^{X'}$ denote the small diagonal of X resp. X' . As explained in [43, Section 6], the argument of [41] gives the equality

$$\gamma \circ \Delta_{sm}^X \circ {}^t(\gamma \times \gamma) = \Delta_{sm}^{X'} \quad \text{in } A^{2n}(X' \times X' \times X').$$

The prescription

$$\Pi_i^{X'} := \gamma \circ \pi_i^X \circ {}^t\gamma \in A^n(X' \times X')$$

defines a CK decomposition for X' . (The $\Pi_i^{X'}$ are orthogonal idempotents thanks to Rieβ's result that $\gamma \circ {}^t\gamma = \Delta_{X'}$ and ${}^t\gamma \circ \gamma = \Delta_X$ [41].)

To see that this CK decomposition $\{\Pi_i^{X'}\}$ is multiplicative, let us consider integers i, j, k such that $i + j \neq k$. It follows from the above equalities that

$$\begin{aligned} \Pi_k^{X'} \circ \Delta_{sm}^{X'} \circ (\Pi_i^{X'} \times \Pi_j^{X'}) &= \gamma \circ \Pi_k^X \circ {}^t\gamma \circ \gamma \circ \Delta_{sm}^X \circ {}^t(\gamma \times \gamma) \circ (\gamma \times \gamma) \circ (\Pi_i^X \times \Pi_j^X) \circ {}^t\gamma \\ &= \gamma \circ \Pi_k^X \circ \Delta_{sm}^X \circ (\Pi_i^X \times \Pi_j^X) \circ {}^t\gamma \\ &= 0 \text{ in } A^{2n}(X' \times X'). \end{aligned}$$

(Here we have again used Rieβ's result that $\gamma \circ {}^t\gamma = \Delta_{X'}$ and ${}^t\gamma \circ \gamma = \Delta_X$.) \square

2.4 – Niveau filtration

DEFINITION 2.14 (Coniveau filtration [10]). Let X be a quasi-projective variety. The *coniveau filtration* on cohomology and on homology is defined as

$$\begin{aligned} N^c H^i(X, \mathbb{Q}) &= \sum \text{Im}(H_Y^i(X, \mathbb{Q}) \rightarrow H^i(X, \mathbb{Q})); \\ N^c H_i(X, \mathbb{Q}) &= \sum \text{Im}(H_i(Z, \mathbb{Q}) \rightarrow H_i(X, \mathbb{Q})), \end{aligned}$$

where Y runs over codimension $\geq c$ subvarieties of X , and Z over dimension $\leq i - c$ subvarieties.

Vial introduced the following variant of the coniveau filtration:

DEFINITION 2.15 (Niveau filtration [48]). Let X be a smooth projective variety. The *niveau filtration* on homology is defined as

$$\widetilde{N}^j H_i(X) = \sum_{\Gamma \in A_{i-j}(Z \times X)} \text{Im}(H_{i-2j}(Z) \rightarrow H_i(X)),$$

where the union runs over all smooth projective varieties Z of dimension $i - 2j$, and all correspondences $\Gamma \in A_{i-j}(Z \times X)$. The niveau filtration on cohomology is defined as

$$\widetilde{N}^c H^i X := \widetilde{N}^{c-i+n} H_{2n-i} X.$$

REMARK 2.16. The niveau filtration is included in the coniveau filtration:

$$\widetilde{N}^j H^i(X) \subset N^j H^i(X).$$

These two filtrations are expected to coincide; indeed, Vial shows this is true if and only if the Lefschetz standard conjecture is true for all varieties [48, Proposition 1.1].

Using the truth of the Lefschetz standard conjecture in degree ≤ 1 , it can be checked [48, page 415 "Properties"] that the two filtrations coincide in a certain range:

$$\widetilde{N}^j H^i(X) = N^j H^i X \text{ for all } j \geq \frac{i-1}{2}.$$

2.5 – Refined CK decomposition

THEOREM 2.17 (Vial [48]). Let X be a smooth projective variety of dimension $n \leq 5$. Assume the Lefschetz standard conjecture $B(X)$ holds (in particular, the Künneth components $\pi_i \in H^{2n}(X \times X)$ are algebraic). Then there is a splitting into mutually orthogonal idempotents

$$\pi_i = \sum_j \pi_{i,j} \in H^{2n}(X \times X),$$

such that

$$(\pi_{i,j})_* H^*(X) = gr_{\mathbb{N}}^j H^i(X) .$$

(Here, the graded $gr_{\mathbb{N}}^j H^i(X)$ can be identified with a Hodge substructure of $H^i(X)$ using the polarization.) In particular,

$$\begin{aligned} (\pi_{2,1})_* H^j(X) &= H^2(X) \cap F^1 , \\ (\pi_{2,0})_* H^j(X) &= H_{tr}^2(X) . \end{aligned}$$

(Here F^* denotes the Hodge filtration, and $H_{tr}^2(X)$ is the orthogonal complement to $H^2(X) \cap F^1$ under the pairing

$$\begin{aligned} H^2(X) \otimes H^2(X) &\rightarrow \mathbb{Q} , \\ a \otimes b &\mapsto a \cup h^{n-2} \cup b .) \end{aligned}$$

PROOF. This is [48, Theorem 1]. □

THEOREM 2.18 (Vial [48]). *Let X be as in theorem 2.17. Assume in addition X has finite-dimensional motive. Then there exists a CK decomposition $\Pi_i \in A^n(X \times X)$, and a splitting into mutually orthogonal idempotents*

$$\Pi_i = \sum_j \Pi_{i,j} \in A^n(X \times X) ,$$

such that

$$\Pi_{i,j} = \pi_{i,j} \text{ in } H^{2n}(X \times X) ,$$

and

$$(\Pi_{2i,i})_* A^k(X) = 0 \text{ for all } k \neq i .$$

The motive $h_{i,0}(X) = (X, \Pi_{i,0}, 0) \in \mathcal{M}_{\text{rat}}$ is well-defined up to isomorphism.

PROOF. This is [48, Theorem 2]. The last statement follows from [48, Proposition 1.8] combined with [31, Theorem 7.7.3]. □

REMARK 2.19. In case X is a surface with finite-dimensional motive, there is equality

$$h_{2,0}(X) = t_2(X) \text{ in } \mathcal{M}_{\text{rat}} ,$$

where $t_2(X)$ is the ‘‘transcendental part of the motive’’ constructed for any surface (not necessarily with finite-dimensional motive) in [31].

LEMMA 2.20. *Let X be a smooth projective variety as in theorem 2.18, and assume*

$$\dim H^2(X, \mathcal{O}_X) = 1 .$$

Then the motive

$$h_{2,0}(X) \in \mathcal{M}_{\text{rat}}$$

is indecomposable, i.e. any non-zero submotive $M \subset h_{2,0}(X)$ is equal to $h_{2,0}(X)$.

PROOF. (This kind of argument is well-known, cf. for instance [55, Corollary 3.11] or [39, Corollary 2.10] where this is proven for $K3$ surfaces with finite-dimensional motive.) The idea is that there are no non-zero Hodge substructures strictly contained in $H_{tr}^2(X)$. Since the motive $M \subset h_{2,0}(X)$ defines a Hodge substructure

$$H^*(M) \subset H_{tr}^2(X) ,$$

we must have $H^*(M) = H_{tr}^2(X)$ and thus an equality of homological motives

$$M = h_{2,0}(X) \text{ in } \mathcal{M}_{\text{hom}} .$$

Using finite-dimensionality of X , it follows there is an equality of Chow motives

$$M = h_{2,0}(X) \text{ in } \mathcal{M}_{\text{rat}} .$$

□

LEMMA 2.21. *Let X_1, X_2 be two projective quotient varieties of dimension 4. Assume X_1, X_2 have finite-dimensional motive, verify the Lefschetz standard conjecture and*

$$N_H^1 H^4(X_j) = \widetilde{N}^1 H^4(X_j) \text{ for } j = 1, 2 ,$$

where N_H^* is the Hodge coniveau filtration. Let $\Gamma \in A^4(X_1 \times X_2)$ and $\Psi \in A^4(X_2 \times X_1)$. The following are equivalent:

(i)

$$\Gamma_*: H^{0,4}(X_1) \rightarrow H^{0,4}(X_2)$$

is an isomorphism, with inverse Ψ_* ;

(ii)

$$\Gamma_*: H_{tr}^4(X_1) \rightarrow H_{tr}^4(X_2)$$

is an isomorphism, with inverse Ψ_* ;

(iii)

$$\Gamma: h_{4,0}(X_1) \rightarrow h_{4,0}(X_2) \text{ in } \mathcal{M}_{\text{rat}}$$

is an isomorphism, with inverse Ψ .

PROOF. Assume (i), i.e.

$$\Psi_* \Gamma_* = \text{id}: H^{0,4}(X_1) \rightarrow H^{0,4}(X_1) .$$

Using the hypothesis $N_H^1 = \widetilde{N}^1$, this implies

$$\Psi_* \Gamma_* = \text{id}: H^4(X_1)/\widetilde{N}^1 \rightarrow H^4(X_1)/\widetilde{N}^1 ,$$

and so

$$(2) \quad (\Psi \circ \Gamma \circ \Pi_{4,0}^{X_1})_* = (\Pi_{4,0}^{X_1})_*: H^*(X_1) \rightarrow H^*(X_1) .$$

Considering the action on $H_{tr}^4(X_1)$, this implies

$$\Psi_* \Gamma_* = \text{id}: H_{tr}^4(X_1) \rightarrow H_{tr}^4(X_1) .$$

Switching the roles of X_1 and X_2 , one finds that likewise $\Gamma_* \Psi_* = \text{id}$ on $H_{tr}^4(X_2)$, and so the isomorphism of (ii) is proven.

Next, we note that it formally follows from equality (2) that Ψ is left-inverse to

$$\Gamma: h_{4,0}(X_1) \rightarrow h_{4,0}(X_2) \text{ in } \mathcal{M}_{\text{hom}} .$$

Switching roles of X_1 and X_2 , one finds Ψ is also right-inverse to Γ and so

$$\Gamma: h_{4,0}(X_1) \rightarrow h_{4,0}(X_2) \text{ in } \mathcal{M}_{\text{hom}}$$

is an isomorphism, with inverse Ψ . By finite-dimensionality, the same holds in \mathcal{M}_{rat} , establishing (iii). □

REMARK 2.22. The equality

$$N_H^1 H^4(X_j) = \widetilde{N}^1 H^4(X_j)$$

in the hypothesis of lemma 2.21 is the conjunction of the generalized Hodge conjecture $N_H^1 = N^1$ and Vial's conjecture $N^1 = \widetilde{N}^1$.

2.6 – Symmetrically distinguished cycles on abelian varieties

DEFINITION 2.23 (O’Sullivan [37]). Let A be an abelian variety. Let $a \in A^*(A)$ be a cycle. For $m \geq 0$, let

$$V_m(a) \subset A^*(A^m)$$

denote the \mathbb{Q} -vector space generated by elements

$$p_*\left((p_1)^*(a^{r_1}) \cdot (p_2)^*(a^{r_2}) \cdot \dots \cdot (p_n)^*(a^{r_n})\right) \in A^*(A^m).$$

Here $n \leq m$, and $r_j \in \mathbb{N}$, and $p_i: A^n \rightarrow A$ denotes projection on the i -th factor, and $p: A^n \rightarrow A^m$ is a closed immersion with each component $A^n \rightarrow A$ being either a projection or the composite of a projection with $[-1]: A \rightarrow A$.

The cycle $a \in A^*(A)$ is said to be *symmetrically distinguished* if for every $m \in \mathbb{N}$ the composition

$$V_m(a) \subset A^*(A^m) \rightarrow A^*(A^m)/A_{hom}^*(A^m)$$

is injective.

THEOREM 2.24 (O’Sullivan [37]). *The symmetrically distinguished cycles form a \mathbb{Q} -subalgebra $A_{sym}^*(A) \subset A^*(A)$, and the composition*

$$A_{sym}^*(A) \subset A^*(A) \rightarrow A^*(A)/A_{hom}^*(A)$$

is an isomorphism. Symmetrically distinguished cycles are stable under pushforward and pullback of homomorphisms of abelian varieties.

REMARK 2.25. For discussion and applications of the notion of symmetrically distinguished cycles, in addition to [37] we refer to [43, Section 7], [50], [3], [20].

LEMMA 2.26. *Let A be an abelian variety of dimension g .*

(i) *There exists an MCK decomposition $\{\Pi_i^A\}$ that is self-dual and consists of symmetrically distinguished cycles.*

(ii) *Assume $g \leq 5$, and let $\{\Pi_i^A\}$ be as in (i). There exists a further splitting*

$$\Pi_2^A = \Pi_{2,0}^A + \Pi_{2,1}^A \quad \text{in } A^g(A \times A),$$

where the $\Pi_{2,i}^A$ are symmetrically distinguished and $\Pi_{2,i}^A = \pi_{2,i}^A$ in $H^{2g}(A \times A)$.

PROOF. (i) An explicit formula for $\{\Pi_i^A\}$ is given in [43, Section 7 Formula (45)].

(ii) The point is that $\Pi_{2,1}^A$ is (by construction) a cycle of type

$$\sum_j C_j \times D_j \quad \text{in } A^g(A \times A),$$

where $D_j \subset A$ is a symmetric divisor and $C_j \subset A$ is a curve obtained by intersecting a symmetric divisor with hyperplanes. This implies $\Pi_{2,1}^A$ is symmetrically distinguished. By assumption, Π_2^A is symmetrically distinguished and hence so is $\Pi_{2,0}^A$. \square

2.7 – The very special EPW sextic

This subsection introduces the main actor of this tale: the very symmetric EPW sextic discovered in [16].

DEFINITION 2.27 ([5]). A *hyperkähler variety* is a simply-connected smooth projective variety X such that $H^0(X, \Omega_X^2)$ is spanned by a nowhere degenerate holomorphic 2-form.

THEOREM 2.28 (Donten–Bury et alii [16]). *Let $X \subset \mathbb{P}^5(\mathbb{C})$ be defined by the equation*

$$\begin{aligned} & x_0^6 + x_1^6 + x_2^6 + x_3^6 + x_4^6 + x_5^6 + (x_0^4 x_1^2 + x_0^4 x_2^2 + \cdots + x_4^2 x_5^4) \\ & + (x_0^2 x_1^2 x_2^2 + x_0^2 x_1^2 x_3^2 + \cdots + x_3^2 x_4^2 x_5^2) + x_0 x_1 x_2 x_3 x_4 x_5 = 0. \end{aligned}$$

(Note that the parentheses are symmetric functions in the variables x_0, \dots, x_5 .)

(i) *The hypersurface X is an EPW sextic (in the sense of [18], [35]).*

(ii) *Let S be the K3 surface obtained from a certain Del Pezzo surface in [51], and let $S^{[2]}$ denote the Hilbert scheme of 2 points on S . Then there is a rational map (of degree 2)*

$$\phi: S^{[2]} \dashrightarrow X.$$

There exists a commutative diagram

$$\begin{array}{ccccc} S^{[2]} & \xrightarrow{\text{flops}} & \overline{S^{[2]}} & \rightarrow & X' := E^4/(G') \leftarrow X_0 \\ & \searrow \phi & & & \swarrow g \\ & & X & & \end{array}$$

Here all horizontal arrows are birational maps. E is an elliptic curve and $X' := E^4/(G')$ is a quotient variety, and X_0 is a hyperkähler variety with $b_2(X_0) = 23$ which is a symplectic resolution of X' . The morphism g is a double cover; X is a projective quotient variety $X = E^4/G$ where $G = (G', i)$ with $i^2 \in G'$. The groups G' and G consist of automorphisms that are group homomorphisms.

(iii) *$S^{[2]}$ and X_0 have finite-dimensional motive and a multiplicative CK decomposition.*

PROOF. (i) [16, Proposition 2.6].

(ii) This is a combination of [16, Proposition 1.1] and [16, Sections 5 and 6]. (Caveat: the group that we denote G' is written G in [16].)

(iii) Vinberg’s K3 surface has Picard number 20; as such, it is a Kummer surface and has finite-dimensional motive. This implies (using [13]) that $S^{[2]}$ has finite-dimensional motive. As birational hyperkähler varieties have isomorphic Chow motives [41], X_0 has finite-dimensional motive. The Hilbert scheme $S^{[2]}$ of any K3 surface S has an MCK decomposition [43, Theorem 13.4]. As the isomorphism of [41] is an isomorphism of algebras in the category of Chow motives, X_0 also has an MCK decomposition (lemma 2.13). \square

REMARK 2.29. The singular locus of the very special EPW sextic X consists of 60 planes. Among these 60 planes, there is a subset of 20 planes which form a complete family of pairwise incident planes in $\mathbb{P}^5(\mathbb{C})$ [16]. This is the maximal number of elements in a complete family of pairwise incident planes, and this seems to be the only known example of a complete family of 20 pairwise incident planes.

REMARK 2.30. The variety X_0 is not unique. In [17, Section 6], it is shown there exist 81^{16} symplectic resolutions of $E^4/(G')$ (some of them non-projective). One noteworthy consequence of theorem 2.28 is that the varieties X_0 are of K3^[2] type (this was not a priori clear from [17]).

REMARK 2.31. For a *generic* EPW sextic X , there exists a hyperkähler fourfold X_0 (called a “double EPW sextic”) equipped with an anti-symplectic involution σ_0 such that $X = X_0/(\sigma_0)$ [35, Theorem 1.1 (2)]. For the very special EPW sextic X , I don’t know whether such X_0 exists. (For this, one would need to show that the Lagrangian subspace A defining the very special EPW sextic is in the Zariski open $\mathrm{LG}(\wedge^3 V)^0 \subset \mathrm{LG}(\wedge^3 V)$ defined in [35, page 3].)

3. Some intermediate steps

3.1 – A strong version of the generalized Hodge conjecture

For later use, we record here a proposition, stating that the very special EPW sextic, as well as some related varieties, satisfy the hypothesis of lemma 2.21:

PROPOSITION 3.1. *Let X_0 be any hyperkähler variety as in theorem 2.28 (i.e., X_0 is a symplectic resolution of $E^4/(G')$). Then*

$$N_H^1 H^4(X_0) = \widetilde{N}^1 H^4(X_0) .$$

(Here N_H^* denotes the Hodge coniveau filtration and \widetilde{N}^* denotes the niveau filtration (definition 2.15).)

The same holds for $X' := E^4/(G')$ and for the very special EPW sextic X :

$$\begin{aligned} N_H^1 H^4(X') &= \widetilde{N}^1 H^4(X') , \\ N_H^1 H^4(X) &= \widetilde{N}^1 H^4(X) . \end{aligned}$$

PROOF. The point is that Vinberg's $K3$ surface S has Picard number 20, and so the corresponding statement is easily proven for $S^{[2]}$:

LEMMA 3.2. *Let S be Vinberg's $K3$ surface. Then*

$$N_H^1 H^4(S^{[2]}) = \widetilde{N}^1 H^4(S^{[2]}) .$$

PROOF. Let $\widetilde{S \times S} \rightarrow S \times S$ denote the blow-up of the diagonal. As is well-known, there are isomorphisms of homological motives

$$\begin{aligned} h(S^{[2]}) &\cong h(\widetilde{S \times S})^{\mathfrak{S}_2} , \\ h(\widetilde{S \times S}) &\cong h(S \times S) \oplus h(S)(1) \quad \text{in } \mathcal{M}_{\text{hom}} , \end{aligned}$$

where \mathfrak{S}_2 denotes the symmetric group on 2 elements acting by permutation. It follows there is a correspondence-induced injection

$$H^4(S^{[2]}) \hookrightarrow H^4(S \times S) \oplus H^2(S) .$$

It thus suffices to prove the statement for $S \times S$. Let us write

$$H^2(S) = N \oplus T := NS(S) \oplus H_{tr}^2(S) .$$

We have

$$\begin{aligned} N_H^1 H^4(S \times S) &= H^4(S \times S) \cap F^1 \\ &= H^0(S) \otimes H^4(S) \oplus H^4(S) \otimes H^0(S) \oplus N \otimes N \oplus N \otimes T \oplus T \otimes N \\ &\quad \oplus (T \otimes T) \cap F^1 . \end{aligned}$$

All but the last summand are obviously in \widetilde{N}^1 . As to the last summand, we have that

$$(T \otimes T) \cap F^1 = (T \otimes T) \cap F^2 .$$

Since the Hodge conjecture is true for $S \times S$ (indeed, S is a Kummer surface and the Hodge conjecture is known for powers of abelian surfaces [1, 7.2.2], [2, 8.1(2)]), there is an inclusion

$$(T \otimes T) \cap F^2 \subset N^2 H^4(S \times S) = \widetilde{N}^2 H^4(S \times S) ,$$

and so the lemma is proven. \square

Since birational hyperkähler varieties have isomorphic cohomology rings [25, Corollary 2.7], and the isomorphism (being given by a correspondence) respects Hodge structures, this proves the result for X_0 . Since X_0 dominates X' and X , the result for X' and X follows. Proposition 3.1 is now proven. \square

3.2 – MCK for quotients of abelian varieties

PROPOSITION 3.3. *Let A be an abelian variety of dimension n , and let $G \subset \text{Aut}_{\mathbb{Z}}(A)$ be a finite group of automorphisms of A that are group homomorphisms. The quotient*

$$X = A/G$$

has a self-dual MCK decomposition.

PROOF. A first step is to show there exists a self-dual CK decomposition for X induced by a CK decomposition on A :

CLAIM 3.4. *Let A and X be as in proposition 3.3, and let $p: A \rightarrow X$ denote the quotient morphism. Let $\{\Pi_i^A\}$ be a CK decomposition as in lemma 2.26(i). Then*

$$\Pi_i^X := \frac{1}{d} \Gamma_p \circ \Pi_i^A \circ {}^t \Gamma_p \in A^n(X \times X), \quad i = 0, \dots, 2n$$

defines a self-dual CK decomposition for X .

To prove the claim, we remark that clearly the given Π_i^X lift the Künneth components of X , and their sum is the diagonal of X . We will make use of the following property:

LEMMA 3.5. *Let A be an abelian variety of dimension n , and let $\{\Pi_i^A\}$ be an MCK decomposition as in lemma 2.26(i). For any $g \in \text{Aut}_{\mathbb{Z}}(A)$, we have*

$$\Pi_i^A \circ \Gamma_g = \Gamma_g \circ \Pi_i^A \quad \text{in } A^n(A \times A).$$

PROOF. Because $g_* H^i(A) \subset H^i(A)$, we have a homological equivalence

$$\Pi_i^A \circ \Gamma_g - \Gamma_g \circ \Pi_i^A = 0 \quad \text{in } H^{2n}(A \times A).$$

But the left-hand side is a symmetrically distinguished cycle, and so it is rationally trivial. \square

To see that Π_i^X is idempotent, we note that

$$\begin{aligned} \Pi_i^X \circ \Pi_i^X &= \frac{1}{d^2} \Gamma_p \circ \Pi_i^A \circ {}^t \Gamma_p \circ \Gamma_p \circ \Pi_i^A \circ {}^t \Gamma_p \\ &= \frac{1}{d} \Gamma_p \circ \Pi_i^A \circ \left(\sum_{g \in G} \Gamma_g \right) \circ \Pi_i^A \circ {}^t \Gamma_p \\ &= \frac{1}{d} \Gamma_p \circ \Pi_i^A \circ \Pi_i^A \circ \left(\sum_{g \in G} \Gamma_g \right) \circ {}^t \Gamma_p \\ &= \frac{1}{d} \Gamma_p \circ \Pi_i^A \circ \left(\sum_{g \in G} \Gamma_g \right) \circ {}^t \Gamma_p \\ &= \frac{1}{d} \Gamma_p \circ \Pi_i^A \circ {}^t \Gamma_p \circ \Gamma_p \circ {}^t \Gamma_p \\ &= \frac{1}{d} \Gamma_p \circ \Pi_i^A \circ {}^t \Gamma_p \circ d \Delta_X \\ &= \Gamma_p \circ \Pi_i^A \circ {}^t \Gamma_p = \Pi_i^X \quad \text{in } A^n(X \times X). \end{aligned}$$

(Here, the third equality is an application of lemma 3.5, and the fourth equality is because Π_i^A is idempotent.) The fact that the Π_i^X are mutually orthogonal is proven similarly; one needs to replace $\Pi_i^X \circ \Pi_i^X$ by $\Pi_i^X \circ \Pi_j^X$ in the above argument. This proves claim 3.4.

Now, it only remains to see that the CK decomposition $\{\Pi_i^X\}$ of claim 3.4 is multiplicative.

CLAIM 3.6. *The CK decomposition $\{\Pi_i^X\}$ given by claim 3.4 is an MCK decomposition.*

To prove claim 3.6, let us consider the composition

$$\Pi_k^X \circ \Delta_{sm}^X \circ (\Pi_i^X \times \Pi_j^X) \in A^n(X \times X),$$

where we suppose $i + j \neq k$. There are equalities

$$\begin{aligned} \Pi_k^X \circ \Delta_{sm}^X \circ (\Pi_i^X \times \Pi_j^X) &= \frac{1}{d^3} \Gamma_p \circ \Pi_k^A \circ {}^t\Gamma_p \circ \Delta_{sm}^X \circ \Gamma_{p \times p} \circ (\Pi_i^A \times \Pi_j^A) \circ {}^t\Gamma_{p \times p} \\ &= \frac{1}{d} \Gamma_p \circ \Pi_k^A \circ \Delta_A^G \circ \Delta_{sm}^A \circ (\Delta_A^G \times \Delta_A^G) \circ (\Pi_i^A \times \Pi_j^A) \circ {}^t\Gamma_{p \times p} \\ &= \frac{1}{d} \Gamma_p \circ \Delta_A^G \circ \Pi_k^A \circ \Delta_{sm}^A \circ (\Pi_i^A \times \Pi_j^A) \circ (\Delta_A^G \times \Delta_A^G) \circ {}^t\Gamma_{p \times p} \\ &= 0 \text{ in } A^{2n}(X \times X \times X). \end{aligned}$$

Here, the first equality is by definition of the Π_i^X , the second equality is lemma 3.7 below, the third equality follows from lemma 3.5, and the fourth equality is the fact that $\{\Pi_i^A\}$ is an MCK decomposition for A (lemma 2.26).

LEMMA 3.7. *There is equality*

$$\begin{aligned} {}^t\Gamma_p \circ \Delta_{sm}^X \circ \Gamma_{p \times p} &= \frac{1}{d} \left(\sum_{g \in G} \Gamma_g \right) \circ \Delta_{sm}^A \circ \left(\left(\sum_{g \in G} \Gamma_g \right) \times \left(\sum_{g \in G} \Gamma_g \right) \right) \\ &= d^2 \Delta_A^G \circ \Delta_{sm}^A \circ (\Delta_A^G \times \Delta_A^G) \text{ in } A^{2n}(A \times A \times A). \end{aligned}$$

PROOF. The second equality is just the definition of Δ_A^G . As to the first equality, we first note that

$$\Delta_{sm}^X = \frac{1}{d} (p \times p \times p)_* (\Delta_{sm}^A) = \frac{1}{d} \Gamma_p \circ \Delta_{sm}^A \circ {}^t\Gamma_{p \times p} \text{ in } A^{2n}(X \times X \times X).$$

This implies that

$${}^t\Gamma_p \circ \Delta_{sm}^X \circ \Gamma_{p \times p} = \frac{1}{d} {}^t\Gamma_p \circ \Gamma_p \circ \Delta_{sm}^A \circ {}^t\Gamma_{p \times p} \circ \Gamma_{p \times p}.$$

But ${}^t\Gamma_p \circ \Gamma_p = \sum_{g \in G} \Gamma_g$, and thus

$${}^t\Gamma_p \circ \Delta_{sm}^X \circ \Gamma_{p \times p} = \frac{1}{d} \left(\sum_{g \in G} \Gamma_g \right) \circ \Delta_{sm}^A \circ \left(\left(\sum_{g \in G} \Gamma_g \right) \times \left(\sum_{g \in G} \Gamma_g \right) \right) \text{ in } A^{2n}(A \times A \times A),$$

as claimed. □

This ends the proof of proposition 3.3. □

In the set-up of proposition 3.3, one can actually say more about certain pieces $A_{(j)}^i(X)$:

PROPOSITION 3.8. *Let $X = A/G$ be as in proposition 3.3. Assume $n = \dim X \leq 5$ and $H^2(X, \mathcal{O}_X) = 0$. Assume also there exists $X' = A/(G')$ where $G = (G', i)$ with $i^2 \in G'$, and the action of i on $H^2(X', \mathcal{O}_{X'})$ is minus the identity. Then any CK decomposition $\{\Pi_i\}$ of X verifies*

$$\begin{aligned} (\Pi_2)_* A^j(X) &= 0 \text{ for all } j \neq 1, \\ (\Pi_6)_* A^j(X) &= 0 \text{ for all } j \neq 3. \end{aligned}$$

PROOF. It suffices to prove this for one particular CK decomposition, in view of the following lemma:

LEMMA 3.9. *Let $X = A/G$ be as in proposition 3.3. Let $\Pi, \Pi' \in A^n(X \times X)$ be idempotents, and assume $\Pi - \Pi' = 0$ in $H^{2n}(X \times X)$. Then*

$$(\Pi)_* A^i(X) = 0 \iff (\Pi')_* A^i(X) = 0 .$$

PROOF. This follows from [48, Lemma 1.14]. Alternatively, here is a direct proof. Let $p: A \rightarrow X$ denote the quotient morphism, and let $d := |G|$. One defines

$$\begin{aligned} \Pi_A &:= \frac{1}{d} {}^t \Gamma_p \circ \Pi \circ \Gamma_p \in A^n(A \times A) , \\ \Pi'_A &:= \frac{1}{d} {}^t \Gamma_p \circ \Pi' \circ \Gamma_p \in A^n(A \times A) . \end{aligned}$$

It is readily checked Π_A, Π'_A are idempotents, and they are homologically equivalent.

Let us assume $(\Pi)_* A^i(X) = 0$ for a certain i . Then also

$$(\Pi_A)_* p^* A^i(X) = \left(\frac{1}{d} {}^t \Gamma_p \circ \Pi \circ \Gamma_p \circ {}^t \Gamma_p \right)_* A^i(X) = ({}^t \Gamma_p \circ \Pi)_* A^i(X) = 0 .$$

By finite-dimensionality of A , the difference $\Pi_A - \Pi'_A \in A_{hom}^n(A \times A)$ is nilpotent, i.e. there exists $N \in \mathbb{N}$ such that

$$(\Pi_A - \Pi'_A)^{\circ N} = 0 \text{ in } A^n(A \times A) .$$

Upon developing, this implies

$$\Pi'_A = (\Pi'_A)^{\circ N} = Q_1 + \dots + Q_N \text{ in } A^n(A \times A) ,$$

where each Q_j is a composition

$$Q_j = Q_j^1 \circ Q_j^2 \circ \dots \circ Q_j^N ,$$

with $Q_j^k \in \{\Pi_A, \Pi'_A\}$, and at least one Q_j^k is Π_A . Since by assumption $(\Pi_A)_* p^* A^i(X) = 0$, it follows that

$$(Q_j)_* = (\text{something})_* (\Pi_A)_* ((\Pi'_A)^{\circ r})_* = 0: p^* A^i(X) \rightarrow p^* A^i(X) \text{ for all } j .$$

But then also

$$(\Pi'_A)_* p^* A^i(X) = (Q_1 + \dots + Q_N)_* p^* A^i(X) = 0 .$$

□

Now, let us take a projector for A of the form

$$\Pi_2^A = \Pi_{2,0}^A + \Pi_{2,1}^A \in A^n(A \times A) ,$$

where $\Pi_{2,0}^A, \Pi_{2,1}^A$ are as in lemma 2.26.

LEMMA 3.10. *Let A be an abelian variety of dimension $n \leq 5$, and let $G \subset \text{Aut}_{\mathbb{Z}}(A)$ be a finite subgroup. Let $\Pi_{2,0}^A$ be as in lemma 2.26. Then*

$$\Pi_{2,0}^A \circ \Delta_A^G = \Delta_A^G \circ \Pi_{2,0}^A \in A^n(A \times A)$$

is idempotent. (Here, as before, we write $\Delta_A^G := \frac{1}{|G|} \sum_{g \in G} \Gamma_g \in A^n(A \times A)$.)

PROOF. For any $g \in G$, we have the commutativity

$$\Pi_{2,0}^A \circ \Gamma_g = \Gamma_g \circ \Pi_{2,0}^A \text{ in } A^n(A \times A) , \text{ for all } g \in G ,$$

established in lemma 2.26(ii). (Indeed, these cycles are symmetrically distinguished by lemma 2.26(ii), and their difference is homologically trivial because an automorphism $g \in G$ respects the niveau filtration.)

This commutativity clearly implies the equality

$$\Pi_{2,0}^A \circ \Delta_A^G = \Delta_A^G \circ \Pi_{2,0}^A \in A^n(A \times A).$$

To check that $\Pi_{2,0}^A \circ \Delta_A^G$ is idempotent, we note that

$$\Pi_{2,0}^A \circ \Delta_A^G \circ \Pi_{2,0}^A \circ \Delta_A^G = \Pi_{2,0}^A \circ \Pi_{2,0}^A \circ \Delta_A^G \circ \Delta_A^G = \Pi_{2,0}^A \circ \Delta_A^G \text{ in } A^n(A \times A).$$

□

Let us write $G = G' \times \{1, i\}$. Since by assumption, $i_* = -\text{id}$ on $H^{2,0}(X')$, we have equality

$$\frac{1}{2} \left(\Pi_{2,0}^A \circ \Delta_A^{G'} + \Pi_{2,0}^A \circ \Delta_A^{G'} \circ \Gamma_i \right) = 0 \text{ in } H^{2n}(A \times A).$$

On the other hand, the left-hand side is equal to the idempotent $\Pi_{2,0}^A \circ \Delta_A^G$. By finite-dimensionality, it follows that

$$\Pi_{2,0}^A \circ \Delta_A^G = 0 \text{ in } A^n(A \times A).$$

Using Poincaré duality, we also have $i_* = -\text{id}$ on $H^{2,4}(X')$, and so (defining $\Pi_{6,2}^A$ as the transpose of $\Pi_{2,0}^A$) there is also an equality

$$\Pi_{6,2}^A \circ \Delta_A^G = \frac{1}{2} \left(\Pi_{6,2}^A \circ \Delta_A^{G'} + \Pi_{6,2}^A \circ \Delta_A^{G'} \circ \Gamma_i \right) = 0 \text{ in } H^{2n}(A \times A),$$

and hence, by finite-dimensionality

$$\Pi_{6,2}^A \circ \Delta_A^G = 0 \text{ in } A^n(A \times A).$$

Since $\Pi_{2,1}^A$ does not act on $A^j(A)$ for $j \neq 1$ (theorem 2.18), we find in particular that

$$(\Pi_2^A)_* = 0: A^j(A)^G \rightarrow A^j(A)^G \text{ for all } j \neq 1.$$

Likewise, since $\Pi_{6,3}^A = {}^t\Pi_{2,1}^A$ does not act on $A^j(A)$ for $j \neq 3$ (theorem 2.18), we also find that

$$(\Pi_6^A)_* = 0: A^j(A)^G \rightarrow A^j(A)^G \text{ for all } j \neq 3.$$

We now consider the CK decomposition for X defined as in lemma 3.4:

$$\Pi_i^X := \frac{1}{d}\Gamma_p \circ \Pi_i^A \circ {}^t\Gamma_p \in A^n(X \times X).$$

This CK decomposition has the required behaviour:

$$\begin{aligned} (\Pi_2^X)_* A^j(X) &= \left(\frac{1}{d}\Gamma_p \circ \Pi_2^A \circ {}^t\Gamma_p \right)_* A^j(X) \\ &= \left(\frac{1}{d}\Gamma_p \right)_* (\Pi_2^A)_* A^j(X) \\ &= \left(\frac{1}{d}\Gamma_p \right)_* (\Pi_2^A)_* A^j(A)^G = 0 \text{ for all } j \neq 1, \end{aligned}$$

and likewise

$$(\Pi_6^X)_* A^j(X) = 0 \text{ for all } j \neq 3.$$

This proves proposition 3.8. □

For later use, we record here a corollary of the proof of proposition 3.8:

COROLLARY 3.11. *Let A be an abelian variety of dimension $n \leq 5$, and let $\Pi_{2,0}^A, \Pi_{2,1}^A$ be as in lemma 2.26(ii). Let $p: A \rightarrow X = A/G$ be a quotient variety with $G \subset \text{Aut}_{\mathbb{Z}}(A)$. The prescription*

$$\Pi_{2,i}^X := \Gamma_p \circ \Pi_{2,i}^A \circ {}^t\Gamma_p \quad \text{in } A^n(X \times X)$$

defines a decomposition in orthogonal idempotents

$$\Pi_2^X = \Pi_{2,0}^X + \Pi_{2,1}^X \quad \text{in } A^n(X \times X).$$

The $\Pi_{2,i}^X$ verify the properties of the refined CK decomposition of theorem 2.18.

PROOF. One needs to check the $\Pi_{2,i}^X$ are idempotent and orthogonal. This easily follows from the fact that the $\Pi_{2,i}^A$ commute with Γ_g for $g \in G$ (lemma 3.10). \square

3.3 – A surjectivity statement

PROPOSITION 3.12. *Let X_0 be a hyperkähler fourfold as in theorem 2.28. Let $A_{(*)}^*(X_0)$ be the bi-grading defined by the MCK decomposition. Then the intersection product map*

$$A_{(2)}^2(X_0) \otimes A_{(2)}^2(X_0) \rightarrow A_{(4)}^4(X_0)$$

is surjective.

The same holds for $X^l := E^4/(G^l)$ as in theorem 2.28: X^l has an MCK decomposition, and the intersection product map

$$A_{(2)}^2(X^l) \otimes A_{(2)}^2(X^l) \rightarrow A_{(4)}^4(X^l)$$

is surjective.

PROOF. The result of Rieß [41] implies there is an isomorphism of bigraded rings

$$A_{(*)}^*(S^{[2]}) \xrightarrow{\cong} A_{(*)}^*(X_0).$$

For the Hilbert scheme of any K3 surface S , the intersection product map

$$A_{(2)}^2(S^{[2]}) \otimes A_{(2)}^2(S^{[2]}) \rightarrow A_{(4)}^4(S^{[2]})$$

is known to be surjective [43, Theorem 3]. This proves the first statement.

For the second statement, the existence of an MCK decomposition for X^l is a special case of proposition 3.3. To prove the surjectivity statement for X^l , we note that $\phi: X_0 \rightarrow X^l$ is a symplectic resolution and so there are isomorphisms

$$\phi^*: H^{p,0}(X^l) \xrightarrow{\cong} H^{p,0}(X_0) \quad (p = 2, 4).$$

Using lemma 2.21 (which is possible thanks to proposition 3.1), this implies there are isomorphisms

$$\phi^*: H_{tr}^p(X^l) \xrightarrow{\cong} H_{tr}^p(X_0) \quad (p = 2, 4).$$

This means there is an isomorphism of homological motives

$${}^t\Gamma_\phi: h_{p,0}(X^l) \xrightarrow{\cong} h_{p,0}(X_0) \quad \text{in } \mathcal{M}_{\text{hom}} \quad (p = 2, 4).$$

By finite-dimensionality, there are isomorphisms of Chow motives

$${}^t\Gamma_\phi: h_{p,0}(X^l) \xrightarrow{\cong} h_{p,0}(X_0) \quad \text{in } \mathcal{M}_{\text{rat}} \quad (p = 2, 4).$$

Taking Chow groups, this implies there are isomorphisms

$$(3) \quad (\Pi_p^{X_0} \circ {}^t\Gamma_\phi \circ \Pi_p^{X^l})_*: (\Pi_p^{X^l})_* A^i(X^l) \rightarrow (\Pi_p^{X_0})_* A^i(X_0) \quad (p = 2, 4).$$

Let us now consider the diagram

$$\begin{array}{ccc} A_{(2)}^2(X_0) \otimes A_{(2)}^2(X_0) & \rightarrow & A_{(4)}^4(X_0) \\ \uparrow & & \uparrow \\ A^2(X_0) \otimes A^2(X_0) & \rightarrow & A^4(X_0) \\ \uparrow & & \uparrow \\ A_{(2)}^2(X') \otimes A_{(2)}^2(X') & \rightarrow & A_{(4)}^4(X') \end{array}$$

Here, the vertical arrows in the upper square are given by projecting to direct summand; the vertical arrows in the lower square are given by ϕ^* . Since pullback and intersection product commute, the lower square commutes. Since $A_{(*)}^*(X_0)$ is a bigraded ring, the upper square commutes.

The composition of vertical arrows is an isomorphism by (3). The statement for X' now follows from the statement for X_0 . \square

4. Main results

4.1 – Splitting of $A^*(X)$

THEOREM 4.1. *Let X be the very special EPW sextic of theorem 2.28. The Chow ring of X is a bigraded ring*

$$A^*(X) = A_{(*)}^*(X),$$

where

$$\begin{aligned} A^1(X) &= A_{(0)}^1(X) = \mathbb{Q}, \\ A^2(X) &= A_{(0)}^2(X), \\ A^3(X) &= A_{(0)}^3(X) \oplus A_{(2)}^3(X) = \mathbb{Q} \oplus A_{hom}^3(X), \\ A^4(X) &= A_{(0)}^4(X) \oplus A_{(4)}^4(X) = \mathbb{Q} \oplus A_{hom}^4(X). \end{aligned}$$

PROOF. It follows from theorem 2.28 that X is a quotient variety $X = E^4/G$ with $G \subset \text{Aut}_{\mathbb{Z}}(A)$. Moreover, there is another quotient variety $X' = E^4/(G')$ where $G = (G', i)$ and $i^2 \in G'$ and such that i acts on $H^2(X', \mathcal{O}_{X'})$ as $-\text{id}$. Applying proposition 3.3, it follows that X has an MCK decomposition $\{\Pi_i^X\}$. Applying proposition 3.8, it follows that

$$\begin{aligned} (\Pi_2^X)_* A^j(X) &= 0 \quad \text{for all } j \neq 1, \\ (\Pi_6^X)_* A^j(X) &= 0 \quad \text{for all } j \neq 3. \end{aligned}$$

The projectors Π_i^X are 0 for i odd. (Indeed, X has no odd cohomology so the Π_i^X are homologically trivial. Using finite-dimensionality, they are rationally trivial.)

The projectors $\{\Pi_i^X\}$ define a multiplicative bigrading

$$A^*(X) = A_{(*)}^*(X),$$

where $A_{(i)}^j(X) := (\Pi_{2j-i}^X)_* A^j(X)$. The fact that $A_{(i)}^j(X) = 0$ for $i < 0$ follows from the corresponding property for abelian fourfolds [6]. Likewise, the fact that

$$A_{(0)}^j(X) \cap A_{hom}^j(X) = 0 \quad \text{for all } j \geq 3$$

follows from the corresponding property for abelian fourfolds [6]. \square

COROLLARY 4.2. *Let X be the very special EPW sextic. The intersection product maps*

$$\begin{aligned} A^2(X) \otimes A^2(X) &\rightarrow A^4(X), \\ A^2(X) \otimes A^1(X) &\rightarrow A^3(X) \end{aligned}$$

have image of dimension 1.

REMARK 4.3. It is instructive to note that for smooth Calabi–Yau hypersurfaces $X \subset \mathbb{P}^{n+1}(\mathbb{C})$, Voisin has proven that the intersection product map

$$A^j(X) \otimes A^{n-j}(X) \rightarrow A^n(X)$$

has image of dimension 1, for any $0 < j < n$ [54, Theorem 3.4], [56, Theorem 5.25] (cf. also [19] for a generalization to generic complete intersections).

In particular, the first statement of corollary 4.2 holds for any smooth sextic in $\mathbb{P}^5(\mathbb{C})$. The second statement of corollary 4.2, however, is not known (and maybe not true) for a general sextic in $\mathbb{P}^5(\mathbb{C})$. It might be that the second statement is specific to EPW sextics, and related to the presence of a hyperkähler fourfold X_0 which is generically a double cover.

REMARK 4.4. Let F^* be the filtration on $A^*(X)$ defined as

$$F^i A^j(X) = \bigoplus_{\ell \geq i} A_{(\ell)}^j(X).$$

For this filtration to be of Bloch–Beilinson type, it remains to prove that

$$F^1 A^2(X) \stackrel{??}{=} A_{hom}^2(X).$$

This would imply the vanishing $A_{hom}^2(X) = 0$ (i.e. the truth of conjecture 1.5 for X).

Unfortunately, we cannot prove this. At least, it follows from the above description that the conjectural vanishing $A_{hom}^2(X) = 0$ would follow from the truth of Beauville’s conjecture

$$A_{hom}^2(E^4) \stackrel{??}{=} A_{(1)}^2(E^4) \oplus A_{(2)}^2(E^4),$$

where E is an elliptic curve.

4.2 – Splitting of $A^*(X^r)$

DEFINITION 4.5. Let X be a projective quotient variety. For any $r \in \mathbb{N}$, and any $1 \leq i < j < k \leq r$, let

$$\begin{aligned} p_j &: X^r \rightarrow X, \\ p_{ij} &: X^r \rightarrow X \times X, \\ p_{ijk} &: X^r \rightarrow X \times X \times X \end{aligned}$$

denote projection on the j -th factor, resp. projection on the i -th and j -th factor, resp. projection on the i -th and j -th and k -th factor.

We define

$$E^*(X^r) \subset A^*(X^r)$$

as the \mathbb{Q} -subalgebra generated by $(p_j)^* A^1(X)$ and $(p_j)^* A^2(X)$ and $(p_{ij})^*(\Delta_X) \in A^4(X^r)$ and $(p_{ijk})^*(\Delta_{sm}^X) \in A^8(X^r)$.

As explained in the introduction, the hypothesis that EPW sextics that are quotient varieties are in the class \mathcal{C} leads to the following concrete conjecture:

CONJECTURE 4.6. *Let $X \subset \mathbb{P}^5(\mathbb{C})$ be an EPW sextic which is a projective quotient variety. Let $r \in \mathbb{N}$. The restriction of the cycle class map*

$$E^i(X^r) \rightarrow H^{2i}(X^r)$$

is injective for all i .

For the very special EPW sextic, we can prove conjecture 4.6 for 0–cycles and 1–cycles:

THEOREM 4.7. *Let X be the very special EPW sextic of definition 2.28. Let $r \in \mathbb{N}$. The restriction of the cycle class map*

$$E^i(X^r) \rightarrow H^{2i}(X^r)$$

is injective for $i \geq 4r - 1$.

PROOF. The product X^r has an MCK decomposition (since X has one, and the property of having an MCK decomposition is stable under taking products [43, Theorem 8.6]). Therefore, there is a bigrading on the Chow ring of X^r . As we have seen (theorem 4.1), $A^1(X) = A_{(0)}^1(X)$ and $A^2(X) = A_{(0)}^2(X)$. Also, it is readily checked that

$$\Delta_X \in A_{(0)}^4(X \times X).$$

(Indeed, this follows from the fact that

$$\Delta_X = \sum_{i=0}^8 \Pi_i^X = \sum_{i=0}^8 \Pi_i^X \circ \Delta_X \circ \Pi_i^X = \sum_{i=0}^8 (\Pi_i^X \times \Pi_{8-i}^X)_* \Delta_X \quad \text{in } A^4(X \times X),$$

where we have used the fact that the CK decomposition is self–dual.) The fact that X has an MCK decomposition implies that

$$\Delta_{sm}^X \in A_{(0)}^8(X \times X \times X)$$

[43, Proposition 8.4].

Clearly, the pullbacks under the projections p_i, p_{ij}, p_{ijk} respect the bigrading. (Indeed, suppose $a \in A_{(0)}^\ell(X)$, which means $a = (\Pi_{2\ell}^X)_*(a)$. Then the pullback $(p_i)^*(a)$ can be written as

$$X \times \cdots \times X \times (\Pi_{2\ell}^X)_*(a) \times X \times \cdots \times X \in A^\ell(X^r),$$

which is the same as

$$(\Pi_0^X \times \cdots \times \Pi_0^X \times \Pi_{2\ell}^X \times \Pi_0^X \times \cdots \times \Pi_0^X)_*(X \times \cdots \times X \times a \times X \times \cdots \times X).$$

This implies that

$$(p_i)^*(a) \in (\Pi_{2\ell}^{X^r})_* A^\ell(X^r) = A_{(0)}^\ell(X^r),$$

where $\Pi_*^{X^r}$ is the product CK decomposition. Another way to prove the fact that the projections p_i, p_{ij}, p_{ijk} respect the bigrading is by invoking [44, Corollary 1.6].)

It follows there is an inclusion

$$E^*(X^r) \subset A_{(0)}^*(X^r).$$

The finite morphism $p^{X^r}: A^r \rightarrow X^r$ induces a split injection

$$(p^{X^r})^*: A_{(0)}^i(X^r) \cap A_{hom}^i(X^r) \rightarrow A_{(0)}^i(A^r) \cap A_{hom}^i(A^r) \quad \text{for all } i.$$

But the right–hand side is known to be 0 for $i \geq 4r - 1$ [6], and so

$$E^i(X^r) \cap A_{hom}^i(X^r) \subset A_{(0)}^i(X^r) \cap A_{hom}^i(X^r) = 0 \quad \text{for all } i \geq 4r - 1.$$

□

REMARK 4.8. As is clear from the proof of theorem 4.7, there is a link with Beauville’s conjectures for abelian varieties: let E be an elliptic curve, and suppose one knows that

$$A_{(0)}^i(E^{4r}) \cap A_{hom}^i(E^{4r}) = 0 \quad \text{for all } i \text{ and all } r.$$

Then conjecture 4.6 is true for the very special EPW sextic.

4.3 – Relation with some hyperkähler fourfolds

THEOREM 4.9. *Let X be the very special EPW sextic of definition 2.28. Let X_0 be one of the hyperkähler fourfolds of [17, Corollary 6.4], and let $f: X_0 \rightarrow X$ be the generically 2 : 1 morphism constructed in [16]. Then X_0 has an MCK decomposition, and there is an isomorphism*

$$f^*: A_{hom}^4(X) \xrightarrow{\cong} A_{(4)}^4(X_0).$$

PROOF. The MCK decomposition for X_0 was established in theorem 2.28. The morphism $f: X_0 \rightarrow X$ of [16] is constructed as a composition

$$f: X_0 \xrightarrow{\phi} X' := E^4/(G') \xrightarrow{g} X,$$

where ϕ is a symplectic resolution and g is the double cover associated to an anti-symplectic involution. This implies f induces an isomorphism

$$f^*: H^{4,0}(X) \xrightarrow{\cong} H^{4,0}(X') \xrightarrow{\cong} H^{4,0}(X_0).$$

In view of the strong form of the generalized Hodge conjecture (proposition 3.1), X_0 and X' and X verify the hypotheses of lemma 2.21. Applying lemma 2.21, we find isomorphisms of Chow motives

$${}^t\Gamma_f: h_{4,0}(X) \xrightarrow{\cong} h_{4,0}(X') \xrightarrow{\cong} h_{4,0}(X_0) \text{ in } \mathcal{M}_{\text{rat}}.$$

Since $(\Pi_{4,i}^X)_* A^4(X) = 0$ for $i \geq 1$ for dimension reasons, we have

$$(\Pi_4^X)_* A^4(X) = (\Pi_{4,0}^X)_* A^4(X),$$

and the same goes for X' and X_0 . It follows that

$$f^*: A_{hom}^4(X) = A^4(h_{4,0}(X)) \xrightarrow{\cong} A^4(h_{4,0}(X_0)) =: A_{(4)}^4(X_0).$$

□

As a corollary, we obtain an alternative description of the splitting $A_{(*)}^*(X_0)$ for the hyperkähler fourfolds X_0 :

COROLLARY 4.10. *Let $f: X_0 \rightarrow X$ be as in theorem 4.9. The splitting $A_{(*)}^*(X_0)$ (given by the MCK decomposition of X_0) verifies*

$$\begin{aligned} A^4(X_0) &= A_{(4)}^4(X_0) \oplus A_{(2)}^4(X_0) \oplus A_{(0)}^4(X_0) \\ &= f^* A_{hom}^4(X) \oplus \ker(A^4(X_0) \xrightarrow{f^*} A^4(X)) \oplus \mathbb{Q}; \\ A^3(X_0) &= A_{(2)}^3(X_0) \oplus A_{(0)}^3(X_0) \\ &= A_{hom}^3(X_0) \oplus H^{3,3}(X_0); \\ A^2(X_0) &= A_{(2)}^2(X_0) \oplus A_{(0)}^2(X_0) \\ &= \ker(A_{hom}^2(X_0) \xrightarrow{f^*} A^2(X)) \oplus A_{(0)}^2(X_0). \end{aligned}$$

REMARK 4.11. Just as we noted for the EPW sextic X (remark 4.4), for this filtration to be of Bloch–Beilinson type one would need to prove that

$$A_{(0)}^2(X_0) \cap A_{hom}^2(X_0) \stackrel{??}{=} 0,$$

which I cannot prove. This situation is similar to that of the Fano varieties F of lines on a very general cubic fourfold: thanks to work of Shen–Vial [43] there is a multiplicative bigrading $A_{(*)}^*(F)$ which has many good properties and interesting alternative descriptions. The main open problem is to prove that

$$A_{(0)}^2(F) \cap A_{hom}^2(F) \stackrel{??}{=} 0,$$

which doesn't seem to be known for any single F .

REMARK 4.12. Conjecturally, the relations of corollary 4.10 should hold for any double EPW sextic X_0 (with X being the quotient of X_0 under the anti-symplectic involution). However, short of knowing X_0 has finite-dimensional motive (as is the case here, thanks to the presence of the abelian variety E^4), this seems difficult to prove. Note that at least, for a general double EPW sextic X_0 , the relations of corollary 4.10 give a concrete description of a filtration on $A^*(X_0)$ that should be the Bloch–Beilinson filtration.

5. Further results

5.1 – Bloch conjecture

CONJECTURE 5.1 (Bloch [9]). *Let X be a smooth projective variety of dimension n . Let $\Gamma \in A^n(X \times X)$ be a correspondence such that*

$$\Gamma_* = 0: H^{p,0}(X) \rightarrow H^{p,0}(X) \text{ for all } p > 0.$$

Then

$$\Gamma_* = 0: A_{hom}^n(X) \rightarrow A_{hom}^n(X).$$

A weak version of conjecture 5.1 is true for the very special EPW sextic:

PROPOSITION 5.2. *Let X be the very special EPW sextic. Let $\Gamma \in A^4(X \times X)$ be a correspondence such that*

$$\Gamma_* = 0: H^{4,0}(X) \rightarrow H^{4,0}(X).$$

Then there exists $N \in \mathbb{N}$ such that

$$(\Gamma^{\circ N})_* = 0: A_{hom}^4(X) \rightarrow A_{hom}^4(X).$$

PROOF. As is well-known, this follows from the fact that X has finite-dimensional motive; we include a proof for completeness' sake.

By assumption, we have

$$\Gamma_* = 0: H^4(X, \mathbb{C})/F^1 \rightarrow H^4(X, \mathbb{C})/F^1$$

(where F^* is the Hodge filtration). Thanks to the “strong form of the generalized Hodge conjecture” (proposition 3.1), this implies that also

$$\Gamma_* = 0: H^4(X, \mathbb{Q})/\tilde{N}^1 \rightarrow H^4(X, \mathbb{Q})/\tilde{N}^1.$$

Using Vial’s refined CK projectors (theorem 2.18), this means

$$\Gamma \circ \Pi_{4,0}^X = 0 \text{ in } H^8(X \times X),$$

or, equivalently,

$$\Gamma - \sum_{(k,\ell) \neq (4,0)} \Gamma \circ \Pi_{k,\ell}^X = 0 \text{ in } H^8(X \times X).$$

By finite-dimensionality, this implies there exists $N \in \mathbb{N}$ such that

$$\left(\Gamma - \sum_{(k,\ell) \neq (4,0)} \Gamma \circ \Pi_{k,\ell}^X \right)^{\circ N} = 0 \text{ in } A^4(X \times X).$$

Upon developing, this gives an equality

$$(4) \quad \Gamma^{\circ N} = Q_1 + \cdots + Q_N \text{ in } A^4(X \times X),$$

where each Q_j is a composition of correspondences

$$Q_j = Q_j^1 \circ Q_j^2 \circ \cdots \circ Q_j^r \in A^4(X \times X),$$

and for each j , at least one Q_j^i is equal to $\Pi_{k,\ell}^X$ with $(k, \ell) \neq (4, 0)$. Since (for dimension reasons)

$$(\Pi_{k,\ell}^X)_* A_{hom}^4(X) = 0 \quad \text{for all } (k, \ell) \neq (4, 0),$$

it follows that

$$(Q_j)_* A_{hom}^4(X) = 0 \quad \text{for all } j.$$

In view of equality (4), we thus have

$$(\Gamma^{\circ N})_* = 0: A_{hom}^4(X) \rightarrow A_{hom}^4(X).$$

□

For special correspondences, one can do better:

PROPOSITION 5.3. *Let X be the very special EPW sextic. Let $\Gamma \in A^4(X \times X)$ be a correspondence such that*

$$\Gamma^* = 0: H^{4,0}(X) \rightarrow H^{4,0}(X).$$

Assume moreover that Γ can be written as

$$\Gamma = \sum_{i=1}^r c_i \Gamma_{\sigma_i} \quad \text{in } A^4(X \times X),$$

with $c_i \in \mathbb{Q}$ and $\sigma_i \in \text{Aut}(X)$ induced by a G -equivariant automorphism $\sigma_i^E: E^4 \rightarrow E^4$, where $X = E^4/G$ and σ_i^E is a group homomorphism. Then

$$\Gamma^* = 0: A_{hom}^4(X) \rightarrow A_{hom}^4(X).$$

PROOF. Let us write $A = E^4$, and $X' := A/(G')$ for the double cover of X with $\dim H^{2,0}(X') = 1$. The projection $g: X' \rightarrow X$ induces an isomorphism

$$g^*: H^{4,0}(X) \xrightarrow{\cong} H^{4,0}(X'),$$

with inverse given by $\frac{1}{d}g_*$. Let $\sigma_i': X' \rightarrow X'$ ($i = 1, \dots, r$) be the automorphism induced by σ_i^E . For each $i = 1, \dots, r$, there is a commutative diagram

$$\begin{array}{ccc} H^{4,0}(X') & \xrightarrow{(\sigma_i')^*} & H^{4,0}(X') \\ g^* \uparrow & & \downarrow g_* \\ H^{4,0}(X) & \xrightarrow{(\sigma_i)^*} & H^{4,0}(X) \end{array}$$

Defining a correspondence

$$\Gamma' = \sum_{i=1}^r c_i \Gamma_{\sigma_i'} \quad \text{in } A^4(X' \times X'),$$

we thus get a commutative diagram

$$\begin{array}{ccc} H^{4,0}(X') & \xrightarrow{(\Gamma')^*} & H^{4,0}(X') \\ g^* \uparrow & & \downarrow g_* \\ H^{4,0}(X) & \xrightarrow{\Gamma^*} & H^{4,0}(X) \end{array}$$

The assumption on Γ^* thus implies that

$$(\Gamma')^* = 0: H^{4,0}(X') \rightarrow H^{4,0}(X').$$

Since (by construction of X') the cup-product map

$$H^{2,0}(X') \otimes H^{2,0}(X') \rightarrow H^{4,0}(X')$$

is an isomorphism of 1-dimensional \mathbb{C} -vector spaces, we must have that

$$(\Gamma')^* = 0: H^{2,0}(X') \rightarrow H^{2,0}(X').$$

It is readily seen this implies

$$(5) \quad {}^t\Gamma' \circ \Pi_{2,0}^{X'} = 0 \quad \text{in } H^8(X' \times X').$$

Let Γ_A denote the correspondence

$$\Gamma_A := \sum_{i=1}^r c_i \Gamma_{\sigma_i^E} \quad \text{in } A^4(A \times A).$$

Let $p': A \rightarrow X' = A/(G')$ denote the quotient morphism. There are relations

$$(6) \quad \begin{aligned} {}^t\Gamma_{\sigma'} &= \frac{1}{|G'|} \Gamma_{p'} \circ {}^t\Gamma_A \circ {}^t\Gamma_{p'} \quad \text{in } A^4(X' \times X'), \\ \Pi_{2,0}^{X'} &= \frac{1}{|G'|} \Gamma_{p'} \circ \Pi_{2,0}^A \circ {}^t\Gamma_{p'} \quad \text{in } A^4(X' \times X') \end{aligned}$$

(the first relation is by construction of the automorphisms σ_i^E ; the second relation can be taken as definition, cf. corollary 3.11). Plugging in these relations in equality (5), one obtains

$$\Gamma_{p'} \circ {}^t\Gamma_A \circ {}^t\Gamma_{p'} \circ \Gamma_{p'} \circ \Pi_{2,0}^A \circ {}^t\Gamma_{p'} = 0 \quad \text{in } H^8(X' \times X').$$

Composing with ${}^t\Gamma_{p'}$ on the left and $\Gamma_{p'}$ on the right, this implies in particular that

$${}^t\Gamma_{p'} \circ \Gamma_{p'} \circ {}^t\Gamma_A \circ {}^t\Gamma_{p'} \circ \Gamma_{p'} \circ \Pi_{2,0}^A \circ {}^t\Gamma_{p'} \circ \Gamma_{p'} = 0 \quad \text{in } H^8(A \times A).$$

Using the standard relation ${}^t\Gamma_{p'} \circ \Gamma_{p'} = \frac{1}{|G'|} \sum_{g \in G'} \Gamma_g$, this simplifies to

$$\left(\sum_{g \in G'} \Gamma_g \right) \circ {}^t\Gamma_A \circ \left(\sum_{g \in G'} \Gamma_g \right) \circ \Pi_{2,0}^A = 0 \quad \text{in } H^8(A \times A).$$

The left-hand side is a symmetrically distinguished cycle which is homologically trivial, and so it is rationally trivial (theorem 2.24). That is,

$$\left(\sum_{g \in G'} \Gamma_g \right) \circ {}^t\Gamma_A \circ \left(\sum_{g \in G'} \Gamma_g \right) \circ \Pi_{2,0}^A = 0 \quad \text{in } A^4(A \times A),$$

in other words

$${}^t\Gamma_{p'} \circ \Gamma_{p'} \circ {}^t\Gamma_A \circ {}^t\Gamma_{p'} \circ \Gamma_{p'} \circ \Pi_{2,0}^A = 0 \quad \text{in } A^4(A \times A).$$

Now we descend again to X' by composing some more on both sides:

$$\Gamma_{p'} \circ {}^t\Gamma_{p'} \circ \Gamma_{p'} \circ {}^t\Gamma_A \circ {}^t\Gamma_{p'} \circ \Gamma_{p'} \circ \Pi_{2,0}^A \circ {}^t\Gamma_{p'} = 0 \quad \text{in } A^4(X' \times X').$$

Using the relations (6), this shimmers down to

$$({}^t\Gamma') \circ \Pi_{2,0}^{X'} = 0 \quad \text{in } A^4(X' \times X').$$

This implies that

$$(\Gamma')^* = 0: A_{hom}^2(X') \rightarrow A_{hom}^2(X').$$

Since $A_{(4)}^4(X')$ equals the image of the intersection product $A_{hom}^2(X') \otimes A_{hom}^2(X') \rightarrow A^4(X')$ (proposition 3.12), we also have that

$$(\Gamma')^* = 0: A_{(4)}^4(X') \rightarrow A_{(4)}^4(X').$$

The commutative diagram

$$\begin{array}{ccc} A_{(4)}^4(X') & \xrightarrow{(\Gamma')^*} & A_{(4)}^4(X') \\ g^* \uparrow & & \uparrow g^* \\ A_{hom}^4(X) & \xrightarrow{\Gamma^*} & A_{hom}^4(X), \end{array}$$

in which vertical arrows are isomorphisms (proof of theorem 4.9), now implies that

$$\Gamma^* = 0: A_{hom}^4(X) \rightarrow A_{hom}^4(X).$$

□

5.2 – Voisin conjecture

Motivated by the Bloch–Beilinson conjectures, Voisin formulated the following conjecture:

CONJECTURE 5.4 (Voisin [52]). *Let X be a smooth Calabi–Yau variety of dimension n . Let $a, a' \in A_{hom}^n(X)$ be two 0–cycles of degree 0. Then*

$$a \times a' = (-1)^n a' \times a \text{ in } A^{2n}(X \times X).$$

It seems reasonable to expect this conjecture to go through for Calabi–Yau’s that are quotient varieties. In particular, conjecture 5.4 should be true for all EPW sextics that are quotient varieties. We can prove this for the very special EPW sextic:

PROPOSITION 5.5. *Let X be the very special EPW sextic. Let $a, a' \in A_{hom}^4(X)$. Then*

$$a \times a' = a' \times a \text{ in } A^8(X \times X).$$

PROOF. As we have seen, there is a finite morphism $p: A \rightarrow X$, where A is an abelian fourfold and

$$p^*: A_{hom}^4(X) \rightarrow A_{(4)}^4(A) = (\Pi_4^A)_* A^4(A)$$

is a split injection. (The inverse to p^* is given by a multiple of p_* .) Proposition 5.5 now follows from the following fact: any $c, c' \in A_{(4)}^4(A)$ verify

$$c \times c' = c' \times c \text{ in } A^8(A \times A);$$

this is [56, Example 4.40].

□

ACKNOWLEDGEMENTS. *The ideas developed in this note grew into being during the Strasbourg 2014–2015 groupe de travail based on the monograph [56]. Thanks to all the participants of this groupe de travail for a stimulating atmosphere. I am very grateful to Bert van Geemen and to the referee for helpful comments, and to Charles Vial for making me appreciate [37], which is an essential ingredient in this note.*

Many thanks to Yasuyo, Kai and Len for hospitably receiving me in the Schiltigheim Math. Research Institute, where this note was written.

REFERENCES

- [1] S. Abdulali, Filtrations on the cohomology of abelian varieties, in: *The arithmetic and geometry of algebraic cycles, Banff 1998* (B. Brent Gordon et al., eds.), CRM Proceedings and Lecture Notes, American Mathematical Society Providence 2000,
- [2] S. Abdulali, Tate twists of Hodge structures arising from abelian varieties, in: *Recent Advances in Hodge Theory: Period Domains, Algebraic Cycles, and Arithmetic* (M. Kerr et al., eds.), London Mathematical Society Lecture Note Series 427, Cambridge University Press, Cambridge 2016,
- [3] G. Ancona, Décomposition de motifs abéliens, *Manuscripta Math.* 146 (3) (2015), 307—328,
- [4] Y. André, Motifs de dimension finie (d’après S.-I. Kimura, P. O’Sullivan,...), *Séminaire Bourbaki 2003/2004*, Astérisque 299 Exp. No. 929, viii, 115—145,
- [5] A. Beauville, Variétés Kähleriennes dont la première classe de Chern est nulle, *J. Differential Geom.* 18 (4) (1984), 755—782,
- [6] A. Beauville, Sur l’anneau de Chow d’une variété abélienne, *Math. Ann.* 273 (1986), 647—651,
- [7] A. Beauville, On the splitting of the Bloch–Beilinson filtration, in: *Algebraic cycles and motives* (J. Nagel et alii, eds.), London Math. Soc. Lecture Notes 344, Cambridge University Press 2007,
- [8] A. Beauville and C. Voisin, On the Chow ring of a $K3$ surface, *J. Alg. Geom.* 13 (2004), 417—426,
- [9] S. Bloch, *Lectures on algebraic cycles*, Duke Univ. Press Durham 1980,
- [10] S. Bloch and A. Ogus, Gersten’s conjecture and the homology of schemes, *Ann. Sci. Ecole Norm. Sup.* 4 (1974), 181—202,
- [11] S. Bloch and V. Srinivas, Remarks on correspondences and algebraic cycles, *American Journal of Mathematics* Vol. 105, No 5 (1983), 1235—1253,
- [12] M. Brion, Log homogeneous varieties, in: *Actas del XVI Coloquio Latinoamericano de Algebra*, Revista Matemática Iberoamericana, Madrid 2007, arXiv: math/0609669,
- [13] M. de Cataldo and L. Migliorini, The Chow groups and the motive of the Hilbert scheme of points on a surface, *Journal of Algebra* 251 no. 2 (2002), 824—848,
- [14] P. Deligne, La conjecture de Weil pour les surfaces $K3$, *Invent. Math.* 15 (1972), 206—226,
- [15] C. Deninger and J. Murre, Motivic decomposition of abelian schemes and the Fourier transform. *J. Reine u. Angew. Math.* 422 (1991), 201—219,
- [16] M. Donten–Bury, B. van Geemen, G. Kapustka, M. Kapustka and J. Wiśniewski, A very special EPW sextic and two IHS fourfolds, arXiv:1509.06214v2, to appear in *Geometry & Topology*,
- [17] M. Donten–Bury and J. Wiśniewski, On 81 symplectic resolutions of a 4–dimensional quotient by a group of order 32, arXiv:1409.4204,
- [18] D. Eisenbud, S. Popescu and C. Walter, Lagrangian subbundles and codimension 3 subcanonical subschemes, *Duke Math. J.* 107(3) (2001), 427—467,
- [19] L. Fu, Decomposition of small diagonals and Chow rings of hypersurfaces and Calabi–Yau complete intersections, *Advances in Mathematics* (2013), 894—924,
- [20] L. Fu, Beauville–Voisin conjecture for generalized Kummer varieties, *International Mathematics Research Notices* 12 (2015), 3878—3898,
- [21] L. Fu, Z. Tian and C. Vial, Motivic hyperkähler resolution conjecture for generalized Kummer varieties, arXiv:1608.04968,
- [22] W. Fulton, *Intersection theory*, Springer–Verlag Ergebnisse der Mathematik, Berlin Heidelberg New York Tokyo 1984,
- [23] V. Guletskiĭ and C. Pedrini, The Chow motive of the Godeaux surface, in: *Algebraic Geometry, a volume in memory of Paolo Francia* (M. Beltrametti et alii, eds.), Walter de Gruyter, Berlin New York, 2002,
- [24] R. Hartshorne, Equivalence relations on algebraic cycles and subvarieties of small codimension, in: *Algebraic geometry, Arcata 1974*, Proc. Symp. Pure Math. Vol. 29, Amer. Math. Soc., Providence 1975,
- [25] D. Huybrechts, The Kähler cone of a compact hyperkähler manifold, *Math. Ann.* 326 (2003), 499—513,
- [26] F. Ivorra, Finite dimensional motives and applications (following S.-I. Kimura, P. O’Sullivan and others), available from <https://perso.univ-rennes1.fr/florian.ivorra/>

- [27] J. Iyer, Murre’s conjectures and explicit Chow–Künneth projectors for varieties with a nef tangent bundle, *Transactions of the Amer. Math. Soc.* 361 (2008), 1667—1681,
- [28] J. Iyer, Absolute Chow–Künneth decomposition for rational homogeneous bundles and for log homogeneous varieties, *Michigan Math. Journal* Vol.60, 1 (2011), 79—91,
- [29] U. Jannsen, Motivic sheaves and filtrations on Chow groups, in: *Motives* (U. Jannsen et alii, eds.), *Proceedings of Symposia in Pure Mathematics* Vol. 55 (1994), Part 1,
- [30] U. Jannsen, On finite–dimensional motives and Murre’s conjecture, in: *Algebraic cycles and motives* (J. Nagel et alii, eds.), Cambridge University Press, Cambridge 2007,
- [31] B. Kahn, J. P. Murre and C. Pedrini, On the transcendental part of the motive of a surface, in: *Algebraic cycles and motives* (J. Nagel et alii, eds.), Cambridge University Press, Cambridge 2007,
- [32] S. Kimura, Chow groups are finite dimensional, in some sense, *Math. Ann.* 331 (2005), 173—201,
- [33] J. Murre, On a conjectural filtration on the Chow groups of an algebraic variety, parts I and II, *Indag. Math.* 4 (1993), 177—201,
- [34] J. Murre, J. Nagel and C. Peters, *Lectures on the theory of pure motives*, Amer. Math. Soc. University Lecture Series 61, Providence 2013,
- [35] K. O’Grady, Irreducible symplectic 4–folds and Eisenbud–Popescu–Walter sextics, *Duke Math. J.* 134(1) (2006), 99—137,
- [36] K. O’Grady, Double covers of EPW–sextics, *Michigan Math. J.* 62 (2013), 143—184,
- [37] P. O’Sullivan, Algebraic cycles on an abelian variety, *J. f. Reine u. Angew. Math.* 654 (2011), 1—81,
- [38] C. Pedrini, On the finite dimensionality of a $K3$ surface, *Manuscripta Mathematica* 138 (2012), 59—72,
- [39] C. Pedrini, Bloch’s conjecture and valences of correspondences for $K3$ surfaces, arXiv:1510.05832v1,
- [40] C. Pedrini and C. Weibel, Some surfaces of general type for which Bloch’s conjecture holds, to appear in: *Period Domains, Algebraic Cycles, and Arithmetic*, Cambridge Univ. Press, 2015,
- [41] U. Rieß, On the Chow ring of birational irreducible symplectic varieties, *Manuscripta Math.* 145 (2014), 473—501,
- [42] T. Scholl, Classical motives, in: *Motives* (U. Jannsen et alii, eds.), *Proceedings of Symposia in Pure Mathematics* Vol. 55 (1994), Part 1,
- [43] M. Shen and C. Vial, The Fourier transform for certain hyperKähler fourfolds, *Memoirs of the AMS* 240 (2016), no.1139,
- [44] M. Shen and C. Vial, The motive of the Hilbert cube $X^{[3]}$, *Forum Math. Sigma* 4 (2016),
- [45] C. Vial, Algebraic cycles and fibrations, *Documenta Math.* 18 (2013), 1521—1553,
- [46] C. Vial, Projectors on the intermediate algebraic Jacobians, *New York J. Math.* 19 (2013), 793—822,
- [47] C. Vial, Remarks on motives of abelian type, *Tohoku Math. J.* 69 (2017), 195—220,
- [48] C. Vial, Niveau and coniveau filtrations on cohomology groups and Chow groups, *Proceedings of the LMS* 106(2) (2013), 410—444,
- [49] C. Vial, Chow–Künneth decomposition for 3– and 4–folds fibred by varieties with trivial Chow group of zero–cycles, *J. Alg. Geom.* 24 (2015), 51—80,
- [50] C. Vial, On the motive of some hyperkähler varieties, *J. f. Reine u. Angew. Math.* 725 (2017), 235—247,
- [51] E. Vinberg, The two most algebraic $K3$ surfaces, *Math. Ann.* 265 (1) (1983), 1—21,
- [52] C. Voisin, Remarks on zero–cycles of self–products of varieties, in: *Moduli of vector bundles*, *Proceedings of the Taniguchi Congress* (M. Maruyama, ed.), Marcel Dekker New York Basel Hong Kong 1994,
- [53] C. Voisin, On the Chow ring of certain algebraic hyper–Kähler manifolds, *Pure and Applied Math. Quarterly* 4 no 3 (2008), 613—649,
- [54] C. Voisin, Chow rings and decomposition theorems for $K3$ surfaces and Calabi–Yau hypersurfaces, *Geometry & Topology* 16 (2012), 433—473,
- [55] C. Voisin, Bloch’s conjecture for Catanese and Barlow surfaces, *J. Differential Geometry* 97 (2014), 149—175,
- [56] C. Voisin, *Chow Rings, Decomposition of the Diagonal, and the Topology of Families*, Princeton University Press, Princeton and Oxford, 2014,
- [57] Z. Xu, Algebraic cycles on a generalized Kummer variety, arXiv:1506.04297v1,
- [58] Q. Yin, Finite–dimensionality and cycles on powers of $K3$ surfaces, *Comment. Math. Helv.* 90 (2015), 503—511.

Received 22-06-2016; revised 28-11-2016