

## Implications of the index of a fixed point subgroup

ERKAN MURAT TÜRKAN (\*)

ABSTRACT – Let  $G$  be a finite group and  $A \leq \text{Aut}(G)$ . The index  $|G : C_G(A)|$  is called the index of  $A$  in  $G$  and is denoted by  $\text{Ind}_G(A)$ . In this paper, we study the influence of  $\text{Ind}_G(A)$  on the structure of  $G$  and prove that  $[G, A]$  is solvable in case where  $A$  is cyclic,  $\text{Ind}_G(A)$  is squarefree and the orders of  $G$  and  $A$  are coprime. Moreover, for arbitrary  $A \leq \text{Aut}(G)$  whose order is coprime to the order of  $G$ , we show that when  $[G, A]$  is solvable, the Fitting height of  $[G, A]$  is bounded above by the number of primes (counted with multiplicities) dividing  $\text{Ind}_G(A)$  and this bound is best possible.

MATHEMATICS SUBJECT CLASSIFICATION (2010). Primary: 20D45; Secondary: 20D10, 20F16.

KEYWORDS. Index, Fixed point subgroup, Automorphism of a group, Solvable Group, Fitting height.

### 1. Introduction

Throughout this paper, we consider only finite groups. To introduce the notation we use in this paper, let  $G$  be a group and  $x \in G$ . The conjugacy class of  $G$  containing  $x$  is denoted by  $x^G$  and its length is denoted by  $\text{Ind}_G(x)$  which is the index  $|G : C_G(x)|$ . The product of solvable normal subgroups of  $G$  which is the maximal solvable normal subgroup of  $G$  in finite case is denoted by  $S(G)$ . The Fitting subgroup of  $G$  is denoted by  $F(G)$ , the Fitting height of a solvable group  $K$  by  $h(K)$  and the set of prime divisors of order of  $G$  by  $\pi(G)$ .

Arithmetical conditions on the length of conjugacy classes of  $G$  influence nonsimplicity, solvability, supersolvability and nilpotency of  $G$ . There are many results in this problem that can be seen in the historical order in [1].

(\*) *Indirizzo dell'A.*: Çankaya University, Ankara, Turkey – Middle East Technical University, Ankara, Turkey  
E-mail: emturkan@cankaya.edu.tr

Let  $A$  be a subgroup of the automorphism group of  $G$  and let the fixed point subgroup  $\{g \in G \mid \alpha(g) = g \text{ for all } \alpha \in A\}$  of  $A$  in  $G$  be denoted by  $C_G(A)$ . The index of  $C_G(A)$  in  $G$ , denoted by  $\text{Ind}_G(A)$  influences solvability of  $G$  and  $[G, A]$ . There are many results for the case where  $C_G(A)$  is small. One of the most famous paper related to this type of problems is probably Higman's result [2]. There are relatively less papers for the case  $C_G(A)$  is large.

One of the papers joining the first type of problems with the latter case of the second type is Kazarin's work [3]. In 1990, he studied the case where  $A = \langle \alpha \rangle$  and  $\text{Ind}_G(\alpha)$  is a prime power. Namely, he proved the following:

**THEOREM 1.1.** ([3], Corollary 1) *Let  $G$  be a finite group and  $\phi$  one of its automorphisms. If  $C_G(\phi)$  contains a Sylow  $r$ -subgroup for all  $r \in \pi(G) \setminus \{p\}$  then  $\phi$  induces the identity automorphism on  $G/S(G)$ .*

This result led us to investigate the structure of  $[G, \alpha]$  when  $\text{Ind}_G(\alpha)$  is divisible by at least two distinct primes, starting with the case  $\text{Ind}_G(\alpha)$  is squarefree. Although the orders of  $G$  and  $\alpha$  are not necessarily coprime in Kazarin's result, the following example shows the indispensability of the coprimeness condition  $(|G|, |\alpha|) = 1$ , in our case.

**EXAMPLE 1.2.** Let  $G = A_5$  and  $\alpha$  be the inner automorphism of  $G$  induced by the transposition  $(1, 2)$ . Then  $\text{Ind}_G(\alpha) = 10$  but  $[A_5, \alpha] = A_5$  is nonabelian simple.

We prove the following theorem as a result in my thesis work ([9], Theorem 2):

**THEOREM 1.3.** *Let  $G$  be a finite group and  $\alpha$  be an automorphism of  $G$  such that  $(|G|, |\alpha|) = 1$ . If  $\text{Ind}_G(\alpha)$  is squarefree then  $[G, \alpha]$  is solvable.*

In the proof of Theorem 1.3, we use the Classification of the Finite Simple Groups (CFSG) to show non-simplicity of  $[G, \alpha]$ .

One may ask if it is possible to replace the assumption that ' $\text{Ind}_G(\alpha)$  is squarefree' with assumption ' $\text{Ind}_G(\alpha)$  is not divisible by 4'. The following example shows that this is not possible:

**EXAMPLE 1.4.** Let  $G = PSL(3, \mathbb{F}_{3^5})$  and let  $\sigma$  be a field automorphism of order 5. Since  $G$  is a simple group, we have  $[G, \sigma] = G$ . Since  $|G| = 2^4 \cdot 3^{15} \cdot 11^4 \cdot 13 \cdot 61 \cdot 4561$ , we have  $(|G|, |\sigma|) = 1$ . It can be seen that

$C_G(\sigma) = PSL(3, \mathbb{F}_3)$  and hence  $|C_G(\sigma)| = 2^4 \cdot 3^3 \cdot 13$ . It follows that,  $\text{Ind}_G(\sigma) = 3^{12} \cdot 11^4 \cdot 61 \cdot 4561$  is odd but  $[G, \sigma]$  is nonabelian simple.

Another work studying consequences of arithmetical properties of  $\text{Ind}_G(A)$  for given pair  $G, A$  with  $A \leq \text{Aut}(G)$  is due to Parker and Quick [4]. They proved the following:

**THEOREM 1.5.** ([4], *Theorem A*) *Let  $G$  be a finite group and let  $A$  be a group of automorphisms of  $G$  such that the orders of  $G$  and  $A$  are coprime. If  $|G : C_G(A)| \leq n$  then  $|[G, A]| \leq n^{\log_2(n+1)}$ .*

Motivated by this result we investigate the influence of  $\text{Ind}_G(A)$  on the nilpotent height of  $[G, A]$  when  $[G, A]$  is a solvable group and  $(|G|, |A|) = 1$ . Namely, we obtain the following as a result in my thesis work ([9], Theorem 1):

**THEOREM 1.6.** *Let  $G$  be a group and  $A \leq \text{Aut}(G)$  such that  $(|G|, |A|) = 1$  and  $\text{Ind}_G(A) = m$ . If  $[G, A]$  is solvable then the Fitting height of  $[G, A]$  is bounded above by  $\ell(m)$  where  $\ell(m)$  is the number of primes dividing  $m$ , counted with multiplicities.*

The Classification of Finite Simple Groups is not needed in the proof of this theorem. The bound given by Theorem 1.6 is best possible because of the example below:

**EXAMPLE 1.7.** Let  $G$  be the group

$$\langle a, b, c, d \mid a^3 = b^7 = c^3 = d^7 = a^{-1}a^c = a^{-1}a^d = b^{-1}b^c = b^{-1}b^d = b^{-2}b^a = d^{-2}d^c = 1 \rangle$$

and let  $\alpha$  be the involutory automorphism of  $G$  given by  $\alpha(a) = cd^5$ ,  $\alpha(b) = d^2$ ,  $\alpha(c) = ab$  and  $\alpha(d) = b^4$ . We observe by [GAP] that  $|G| = 441$ ,  $|[G, \alpha]| = 147$ ,  $F([G, \alpha]) = 49$ ,  $F_2([G, \alpha]) = [G, \alpha]$ ,  $C_G(\alpha) = \langle abc, b^{a^{-1}}d \rangle$  and  $|G : C_G(\alpha)| = 21$ . Now,  $(|G|, |A|) = (441, 2) = 1$ ,  $\ell([G : C_G(\alpha)]) = \ell(21) = 2$  and Fitting height of  $[G, \alpha]$  is 2.

## 2. The Proof of Theorem 1.3

**PROOF.** We use induction on the order of the semidirect product  $G\langle\alpha\rangle$ .

Suppose  $G$  is a group and  $\alpha \in \text{Aut}(G)$  so that the semidirect product  $G\langle\alpha\rangle$  has the smallest order among all the pairs  $(G, \alpha)$  that satisfies the hypothesis of Theorem 1.3 but  $[G, \alpha]$  is not solvable.

We deduce a contradiction over a series of steps.

Let  $p$  be a prime divisor of order of  $\alpha$ . Then there is a positive integer  $k$  so that  $|\alpha| = kp$ .

Suppose  $k > 1$ . Since  $C_G(\alpha) \leq C_G(\alpha^k)$ , we have  $\text{Ind}_G(\alpha^k)$  divides  $\text{Ind}_G(\alpha)$ . Hence,  $\text{Ind}_G(\alpha^k)$  is squarefree. As  $|G\langle\alpha^k\rangle| < |G\langle\alpha\rangle|$ , we have  $[G, \alpha^k]$  is solvable.

Now,  $[G, \alpha^k]$  is an  $\alpha$ -invariant normal subgroup of  $G$  and so  $\alpha$  induces an automorphism by  $G/[G, \alpha^k]$ . Clearly,  $\text{Ind}_{G/[G, \alpha^k]}(\alpha)$  divides  $\text{Ind}_G(\alpha)$ . So  $\text{Ind}_{G/[G, \alpha^k]}(\alpha)$  is squarefree. It follows by induction assumption that  $[G/[G, \alpha^k], \alpha] = [G, \alpha]/[G, \alpha^k]$  is solvable. Therefore,  $[G, \alpha]$  is solvable which contradicts to our assumption.

Hence,  $k = 1$  and  $\alpha$  is of prime order  $p$ .

Let  $N$  be a proper normal subgroup of  $G\langle\alpha\rangle$ .

Suppose  $\alpha \in N$ . Then  $N = N_1\langle\alpha\rangle$  where  $N_1 = N \cap G$  which is  $\alpha$ -invariant. Since  $\text{Ind}_{N_1}(\alpha)$  divides  $\text{Ind}_G(\alpha)$ , we get  $\text{Ind}_{N_1}(\alpha)$  is squarefree. So by induction assumption  $[N_1, \alpha] = [N, \alpha]$  is solvable. Now,  $\langle\alpha^N\rangle = [N, \alpha]\langle\alpha\rangle$  is solvable. It follows that,  $\alpha \in S(N)$  and as  $S(N) \leq S(G\langle\alpha\rangle)$  we get  $\alpha \in S(G\langle\alpha\rangle)$ . So,  $\langle\alpha^{G\langle\alpha\rangle}\rangle \leq S(G\langle\alpha\rangle)$ . As  $[G, \alpha] \leq \langle\alpha^{G\langle\alpha\rangle}\rangle = \langle\alpha^G\rangle = [G, \alpha]\langle\alpha\rangle$ , we have  $[G, \alpha]$  is solvable, which is a contradiction.

Hence,  $\alpha$  is not contained in a proper normal subgroup of  $G\langle\alpha\rangle$ .

Consider the quotient group  $G\langle\alpha\rangle/N$ .

$\text{Ind}_{G\langle\alpha\rangle/N}(\alpha N)$  is squarefree since it is a divisor of  $\text{Ind}_G(\alpha)$ . By induction,  $[G\langle\alpha\rangle/N, \alpha N]$  is solvable. It follows that,

$$\langle(\alpha N)^{G\langle\alpha\rangle/N}\rangle = [G\langle\alpha\rangle/N, \alpha N]\langle\alpha N\rangle$$

is solvable and hence  $(\alpha N)^{G\langle\alpha\rangle/N} \in S(G\langle\alpha\rangle/N)$ .

Now,  $S(G\langle\alpha\rangle/N) = X/N$  for some normal subgroup  $X$  of  $G\langle\alpha\rangle$  and  $\alpha \in X$ . It follows that,  $X = G\langle\alpha\rangle$ .

Therefore, for any proper normal subgroup  $N$  of  $G\langle\alpha\rangle$ , we have  $G\langle\alpha\rangle/N$  is solvable.

Suppose  $S(G\langle\alpha\rangle) \neq 1$ . Since  $S(G\langle\alpha\rangle)$  is a proper normal subgroup of  $G\langle\alpha\rangle$ , we get  $G\langle\alpha\rangle/S(G\langle\alpha\rangle)$  is solvable. Hence,  $G\langle\alpha\rangle$  is solvable, a contradiction. Therefore,  $S(G\langle\alpha\rangle) = 1$ .

Let  $K$  be a minimal normal subgroup of  $G\langle\alpha\rangle$ . If  $K \not\leq G$ , then  $K \cap G = 1$ . Hence,  $|K| = p$  is prime, which leads the contradiction  $K \leq S(G\langle\alpha\rangle) = 1$ . Thus,  $K \leq G$ .

Suppose  $K \neq G$ . Then  $\text{Ind}_K(\alpha)$  is squarefree since it divides  $\text{Ind}_G(\alpha)$ . Hence, by induction  $[K, \alpha]$  is solvable and so is  $\langle\alpha^K\rangle = [K, \alpha]\langle\alpha\rangle$ . Then

we get  $\alpha \in \langle \alpha^K \rangle = \langle \alpha^{K\langle \alpha \rangle} \rangle \leq S(K\langle \alpha \rangle)$ . Now,

$$[K, \alpha] \leq S(K\langle \alpha \rangle) \cap K \leq S(K) \leq S(G\langle \alpha \rangle) = 1.$$

It follows that,  $\alpha \in C_{G\langle \alpha \rangle}(K) \trianglelefteq G\langle \alpha \rangle$ . So  $C_{G\langle \alpha \rangle}(K) = G\langle \alpha \rangle$ . Then we get the contradiction  $K \leq Z(G\langle \alpha \rangle) \leq S(G\langle \alpha \rangle) = 1$ .

Therefore,  $G$  is the unique minimal normal subgroup of  $G\langle \alpha \rangle$ ,  $G$  is characteristically simple and  $(G\langle \alpha \rangle)' = G$ . Hence,  $G$  is a product of isomorphic copies of a simple group say,  $E \leq G$ . As  $G$  is not solvable,  $E$  is nonabelian.

Suppose  $G \neq E$ . Consider the family  $\{E^{\alpha^k} | k = 0, 1, 2, \dots, p-1\}$  of subgroups of  $G$ . The subgroup  $M = E \times E^\alpha \times \dots \times E_{\alpha^{p-1}}$  is an  $\alpha$ -invariant normal subgroup of  $G$ . So  $M \trianglelefteq G\langle \alpha \rangle$ . Since  $G$  is the unique minimal normal subgroup of  $G\langle \alpha \rangle$  we get  $G = M$ . It follows that  $C_G(\alpha) = \{xx^\alpha x^{\alpha^2} \dots x^{\alpha^{p-1}} | x \in E\}$ . Hence  $|C_G(\alpha)| = |E|$  and  $\text{Ind}_G(\alpha) = |E|^{p-1}$ . Since 2 is a divisor of  $|E|$  and  $(|G|, |\alpha|) = 1$ , we have  $p > 2$  and so  $\text{Ind}_G(\alpha)$  is divisible by  $|E|^2$ , a contradiction.

Therefore,  $G = E$  is a nonabelian simple group.

From Atlas of Finite Groups [8], we observe that  $G$  is not a sporadic simple group as they have no coprime automorphism. Since alternating groups has no coprime automorphism, we get  $G$  is not an alternating group. Thus,  $G$  is a simple group of Lie type. It follows that  $\alpha$  is a field automorphism up to conjugation since  $(|\alpha|, |G|) = 1$ .

Let  $r$  be a prime number. Let  $n_r$  denote the largest power of  $r$  that divides  $n$  and  $L(q)$  denote a simple group of Lie type over the finite field of order  $q$ . By Proposition 4.9.1 in [5], if  $q = r^{ps}$  for some integer  $s$  and  $G = L(q)$  and  $\alpha$  is a field automorphism of order  $p$ , then  $C_G(\alpha) \cong L(r^s)$ .

Let  $G = A_m(r^{ps})$  for  $m \geq 2$  then  $C_G(\alpha) = A_m(r^s)$ . It follows that,

$$\text{Ind}_G(\alpha)_r = (r^{ps})^{m(m+1)/2} / (r^s)^{m(m+1)/2} = r^{sm(m+1)(p-1)/2}$$

For each other family of simple groups of Lie type the argument is the same. In all cases,  $r^2$  divides  $\text{Ind}_G(\alpha)$ . This contradiction completes the proof.  $\square$

### 3. The Proof of Theorem 1.6

PROOF OF THEOREM 1.6. We use induction on the order of  $G$ . Let  $G$  be a minimal counter example to Theorem 1.6 and  $A \leq \text{Aut}(G)$  as in Hypothesis of Theorem 1.6.

Suppose that  $[G, A]$  is properly contained in  $G$ . Since  $(|G|, |A|) = 1$ , we have  $G = [G, A] C_G(A)$  by Lemma 8.2.7 in [6]. It follows that

$$\begin{aligned} |[G, A] : C_{[G, A]}(A)| &= |[G, A] : ([G, A] \cap C_G(A))| \\ &= |[G, A] C_G(A) : C_G(A)| \\ &= |G : C_G(A)| = m \end{aligned}$$

As  $|[G, A]| < |G|$ , by minimality of  $G$ , we get  $h([G, A, A]) \leq \ell(m)$ .

By Lemma 8.2.7 in [6], we know  $[G, A, A] = [G, A]$  since  $(|G|, |A|) = 1$ . This leads to the contradiction  $h([G, A]) \leq \ell(m)$ .

Hence,  $[G, A] = G$ .

If  $G$  is nilpotent then  $h(G) = 1 \leq \ell(m)$ . Thus, we may assume that  $F(G) \not\leq G$ .

Next, suppose that  $F(G)$  is a subgroup of  $C_G(A)$ . As  $[F(G), G] \leq F(G)$ , we have  $[F(G), G, A] = 1$  and  $[A, F(G), G] = 1$ . It follows by the Three Subgroup Lemma (2.2 Theorem 2.3 in [7]) that  $[G, A, F(G)] = [G, F(G)] = 1$ .

Since  $C_G(F(G)) \subseteq F(G)$  by 6.1 Theorem 1.3 in [7], we get  $G = F(G)$ , which is not the case. Hence,  $F(G) \not\leq C_G(A)$ .

Now,  $C_G(A) F(G) \neq C_G(A)$  and hence,

$$\ell(|G : C_G(A) F(G)|) \not\leq \ell(m).$$

As  $|G/F(G)| < |G|$ , we have

$$\begin{aligned} h(G) - 1 &= h(G/F(G)) \\ &\leq \ell(|G/F(G) : C_{G/F(G)}(A)|) \\ &\leq \ell(|G/F(G) : C_G(A) F(G)/F(G)|) \\ &\leq \ell(|G : C_G(A) F(G)|) < \ell(m). \end{aligned}$$

Consequently,  $h([G, A]) - 1 = h(G) - 1 \leq \ell(m) - 1$  and hence  $h([G, A]) \leq \ell(m)$ , completing the proof.  $\square$

#### REFERENCES

- [1] A. R. CAMINA – R. D. CAMINA, *The influence of conjugacy class sizes on the structure of finite groups: A survey*, Asian-Eur. J. Math. **4** (4) (2011), pp. 559–588.
- [2] G. HIGMAN, *Groups and rings having automorphisms without non-trivial fixed elements*, J. London Math. Soc., **32** (1957), pp. 321–334.
- [3] L. S. KAZARIN, *Burnside's  $p^\alpha$  Lemma*, (Russian), Math.Zametki, **48** (1990) no.2, pp. 45–48 ; Translation in Math.Notes, **48** (1991) no:1-2, pp. 749–751

- [4] C. PARKER – M. QUICK, *Coprime automorphisms and their commutators*, J.Algebra, **244** (2001), pp. 260–272.
- [5] D. GORENSTEIN – R. LYONS – R. SOLOMON, *The Classification of the Finite Simple Groups, Number 3*, Mathematical Surveys and Monographs, Vol. 40, Am. Math. Soc., Providence, 1998.
- [6] H. KURZWEIL – B. STELLMACHER, *The Theory of Finite Groups An Introduction*, Springer, New York, 2004.
- [7] D. GORENSTEIN, *Finite Groups*, AMS Chelsea Publishing, New York, 2007.
- [8] J. H. CONWAY – R. T. CURTIS – S. P. NORTON – R. A. PARKER – R. A. WILSON, *Atlas of Finite Groups*, Oxford Univ. Press, Oxford, 1985.
- [9] E. M. TÜRKAN *On the Index of Fixed Point Subgroup*, Ph. D. Thesis, Middle East Technical University, Ankara, 2011
- [GAP] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.8.10*; 2018, (<https://www.gap-system.org>).