

A note on S -semipermutable subgroups of finite groups

HAORAN YU*

ABSTRACT – In this note, we obtain some criteria for p -supersolvability of a finite group and extend some known results concerning weakly S -semipermutable subgroups.

MATHEMATICS SUBJECT CLASSIFICATION (2010). 20D10.

KEYWORDS. p -supersolvability, weakly S -semipermutable subgroups.

1. Introduction

Throughout the paper, we suppose G is a finite group and p is a prime. Let $\pi(G)$ be the set of all the prime divisors of $|G|$. Let $O^p(G) = \bigcap \{N \mid N \trianglelefteq G \text{ and } G/N \text{ is a } p\text{-group}\}$. To state our results, we need to recall some notation. According to Kegel (see [6]), let H be a subgroup of a finite group G ; then H is called an S -permutable subgroup of G if H permutes with every Sylow subgroup of G . According to Chen (see [2]), let H be a subgroup of a finite group G ; then H is said to be S -semipermutable in G if $HQ = QH$ for all Sylow q -subgroups Q of G for all primes q not dividing $|H|$. According to Li et al. (see [7]), let H be a subgroup of a finite group G ; then H is called a weakly S -semipermutable subgroup of G if there exist $T \trianglelefteq \trianglelefteq G$ and $H_1 \leq G$ such that $G = HT$, $H \cap T \leq H_1 \leq H$ and H_1 is S -semipermutable in G . Following Yakov Berkovich and I. M. Isaacs (see [1]), if G is a finite group and p is a prime divisor of $|G|$, we write G_p^* to denote the unique smallest normal subgroup of G for which the corresponding factor group is abelian of exponent dividing $p - 1$. It is well known that G is p -supersolvable if and only if G_p^* is p -nilpotent (see Lemma 3.6 of [1]).

Recently, we proved the following theorem.

*The author is grateful to the referee who provided his/her valuable suggestions and detailed report.

THEOREM 1.1 (Theorem 1.2, [9]). *Let p be a prime dividing the order of a finite group G , e be a positive integer and $P \in \text{Syl}_p(G)$ with $|P| \geq p^{e+1}$. Then G is p -supersolvable if and only if $|P \cap O^p(G_p^*)| \leq p^e$ and $P_1 \cap O^p(G_p^*)$ is S -permutable in G for all subgroups $P_1 \trianglelefteq P$ with $|P_1| = p^e$.*

In this note, at first, we generalize Theorem 1.2 of [9] as follows.

THEOREM 1.2. *Let p be a prime dividing the order of a finite group G , e be a positive integer, $P \in \text{Syl}_p(G)$ with $|P| \geq p^{e+1}$ and $L \trianglelefteq G$ with $O^p(G_p^*) \leq L \leq G$. Suppose that $|P \cap L| \leq p^e$ and $P_1 \cap L$ is S -semipermutable in G for all subgroups $P_1 \trianglelefteq P$ with $|P_1| = p^e$. Then G is p -supersolvable.*

Using Theorem 1.2, we prove the following results which generalize Theorem 1.3 and Theorem 1.4 of [9].

THEOREM 1.3. *Let p be a prime dividing the order of a finite group G , $e \geq 2$ be an integer, $P \in \text{Syl}_p(G)$ with $|P| \geq p^{e+1}$ and $L \trianglelefteq G$ with $O^p(G_p^*) \leq L \leq G$. Suppose that $P_1 \cap L$ is S -semipermutable in G for all subgroups $P_1 \leq P$ with $|P_1| = p^e$. Then G is p -supersolvable.*

Let p be a prime and P be a nonidentity p -group with $|P| = p^n$. We define the set $\Omega(P)$. If $p = 2$ and P is non-abelian, let $\Omega(P) = \{P_1 \mid P_1 \leq P \text{ and } |P_1| = 2\} \cup \{P_2 \mid P_2 \leq P \text{ and } P_2 \text{ is a cyclic subgroup of order } 4\}$. Otherwise, let $\Omega(P) = \{P_1 \mid P_1 \leq P \text{ and } |P_1| = p\}$.

THEOREM 1.4. *Let p be a prime dividing the order of a finite group G , $P \in \text{Syl}_p(G)$ and $L \trianglelefteq G$ with $O^p(G_p^*) \leq L \leq G$. Suppose that $P_1 \cap L$ is S -semipermutable in G for all subgroups $P_1 \in \Omega(P)$. Then G is p -supersolvable.*

Note that Theorem 1.3 and Theorem 1.4 also generalize Theorem 3.5 of [7] and Theorem 3.1, Theorem 3.4 of [8].

2. Preliminaries

LEMMA 2.1. *Let p be a prime dividing the order of a finite group G . Then the following results hold.*

(a) (Lemma 3.1, [1]) *Let $P_1 \leq G$ be a p -group and $N \trianglelefteq G$. If P_1 is S -semipermutable in G , then $P_1 N / N$ is S -semipermutable in G / N .*

(b) (Lemma 3.2, [1]) *Let $P_1 \leq G$ be a p -group and N be a normal p -subgroup of G . If P_1 is S -semipermutable in G , then $P_1 \cap N$ is normalized by $O^p(G)$. In particular, if $P_1 \leq N$, then P_1 is normalized by $O^p(G)$.*

(c) (Lemma 3.3, [1]) *Let $X \leq H \leq G$. If X is S -semipermutable in G , then X is S -semipermutable in H .*

LEMMA 2.2 (Lemma 2.1, [9]). *Let p be a prime dividing the order of a finite group G , $P \in \text{Syl}_p(G)$, $N \trianglelefteq G$ and let e be a positive integer. Write $P_1 = P \cap N$. Assume that $P_1 \trianglelefteq N$ and N is not p -nilpotent. Also assume that $|P_1| \leq p^e$ and $|P| \geq p^{e+1}$. Then P has a normal subgroup P_2 of order p^e with $[P_1 : P_1 \cap P_2] = p$.*

LEMMA 2.3 (Lemma 2.2, [9]). *Let p be a prime dividing the order of a finite group G and $P \in \text{Syl}_p(G)$. Write $\hat{P} = P \cap O^p(G_p^*)$. Assume that $\hat{P} > 1$ and \hat{P} has a maximal subgroup T with $T \trianglelefteq G$. Then $\hat{P} \not\trianglelefteq G$.*

LEMMA 2.4 (Lemma 2.8, [9]). *Let p be a prime dividing the order of a finite group G and P_1 be a p -subgroup of G . Let $L \trianglelefteq G$ and N be a normal p' -subgroup of G . Then $P_1N/N \cap LN/N = (P_1 \cap L)N/N$.*

LEMMA 2.5 (Lemma 2.9, [9]). *Let p be a prime dividing the order of a finite group G and $N \trianglelefteq G$. Then $(G/N)_p^* = G_p^*N/N$, $O^p(G/N) = O^p(G)N/N$ and $O^p((G/N)_p^*) = O^p(G_p^*)N/N$.*

Recently, I. M. Isaacs proved the following significant theorem.

LEMMA 2.6 (Theorem A, [5]). *Let p be a prime dividing the order of a finite group G and P_1 be an S -semipermutable p -subgroup of G . Then P_1^G is solvable.*

Recently, Yakov Berkovich and I. M. Isaacs proved the following powerful results.

LEMMA 2.7 (Yakov Berkovich and I. M. Isaacs). *Let p be a prime and P be a nonidentity finite p -group. Let A act on P via automorphisms.*

(a) (Lemma 2.1(a), [1]) *If P is cyclic, then $O^p(A_p^*)$ acts trivially on P .*

(b) (Theorem A, [1]) *Fix an integer $e \geq 3$. If P is a noncyclic p -group with $|P| \geq p^{e+1}$ and every noncyclic subgroup of P with order p^e is stabilized by $O^p(A)$, then $O^p(A_p^*)$ acts trivially on P .*

(c) (Corollary B, [1]) *If P is a noncyclic p -group with $|P| \geq p^3$ and every subgroup of P with order p^2 is stabilized by $O^p(A)$, then $O^p(A_p^*)$ acts trivially on P .*

LEMMA 2.8 (Yakov Berkovich and I. M. Isaacs). *Let p be a prime dividing the order of a finite group G and $P \in \text{Syl}_p(G)$.*

(a) (Lemma 3.8, [1]) *If P is cyclic and some nonidentity subgroup $U \leq P$ is S -semipermutable in G , then G is p -supersolvable.*

(b) (Theorem D, [1]) *Fix an integer $e \geq 3$. If P is a noncyclic p -group with $|P| \geq p^{e+1}$ and every noncyclic subgroup of P with order p^e is S -semipermutable in G , then G is p -supersolvable.*

(c) (Corollary E, [1]) *If P is a noncyclic p -group with $|P| \geq p^3$ and every subgroup of P with order p^2 is S -semipermutable in G , then G is p -supersolvable.*

LEMMA 2.9 (Lemma 2.12, [9]). *Let p be a prime and P be a nonidentity finite p -group. Let A act on P via automorphisms. Assume that for all $P_1 \in \Omega(P)$, P_1 is stabilized by $O^p(A)$. Then $O^p(A_p^*)$ acts trivially on P .*

3. Main Results

PROOF OF THEOREM 1.2. Suppose that G is a counterexample with minimal order; we complete the following steps to obtain a contradiction. Since G is not p -supersolvable, it follows that $O^p(G_p^*)$ is not p -nilpotent.

Step 1 $P \cap L \geq P \cap O^p(G_p^*) > 1$.

Since $O^p(G_p^*)$ is not p -nilpotent, it follows that $P \cap O^p(G_p^*) > 1$. Since $L \geq O^p(G_p^*)$, it follows that $P \cap L \geq P \cap O^p(G_p^*) > 1$.

Step 2 $O_{p'}(G) = 1$.

By Lemma 2.1(a), Lemma 2.4 and Lemma 2.5, the hypotheses are inherited by $G/O_{p'}(G)$. If $O_{p'}(G) > 1$, then $G/O_{p'}(G)$ is p -supersolvable, and thus G is p -supersolvable. This is a contradiction. Hence $O_{p'}(G) = 1$.

Step 3 Let $\hat{P} = P \cap O^p(G_p^*)$. Then $\hat{P} \trianglelefteq G$.

Let $U = P \cap L$. Then $U \trianglelefteq P$ and $|U| \leq p^e$. Hence P has a normal subgroup P_1 of order p^e with $U \leq P_1$. By the hypotheses, $U = P_1 \cap L$ is S -semipermutable in G . By Lemma 2.6, U^G is solvable. By Step 1, $U^G \geq U > 1$. By Step 2, $O_{p'}(U^G) = 1$. Since U^G is solvable, it follows that $O_p(U^G) > 1$. Let $N = O_p(U^G)$. Since $U^G \leq L$, it follows that $N \leq U$. Hence $|N| \leq |U| \leq p^e$.

Assume that $1 < |N| < p^e$. Recall that $N \leq P \cap L$. By Lemma 2.1(a) and Lemma 2.5, the hypotheses are inherited by G/N . Hence G/N is p -supersolvable. By Lemma 2.5, it follows that $O^p(G_p^*)N/N = O^p((G/N)_p^*)$ is a p' -group, and thus $N \cap O^p(G_p^*)$ is the normal Sylow p -subgroup of $O^p(G_p^*)$. Hence $\hat{P} = N \cap O^p(G_p^*)$, and thus $\hat{P} \trianglelefteq G$.

Assume that $|N| = p^e$. Since $N \leq U$ and $|U| \leq p^e$, it follows that $U = N$ is a normal subgroup of G . Hence $\hat{P} = P \cap O^p(G_p^*) = U \cap O^p(G_p^*) \trianglelefteq G$.

Step 4 The final contradiction.

Recall that \hat{P} is the normal Sylow p -subgroup of $O^p(G_p^*)$ (Step 3), $O^p(G_p^*)$ is not p -nilpotent, $|P| \geq p^{e+1}$ and $1 < |\hat{P}| \leq |U| \leq p^e$. By Lemma 2.2, P has a normal subgroup P_2 of order p^e with $[\hat{P} : \hat{P} \cap P_2] = p$. Note that $\hat{P} \cap P_2 \trianglelefteq P$. By the hypotheses, $P_2 \cap L$ is S -semipermutable in G . By Lemma 2.1(b), $\hat{P} \cap P_2 = \hat{P} \cap P_2 \cap L$ is normalized by $O^p(G)$. Hence $\hat{P} \cap P_2 \trianglelefteq G$. By Lemma 2.3, $\hat{P} \not\trianglelefteq G$. This is a contradiction since $\hat{P} \trianglelefteq G$. Hence we obtain the final contradiction. \square

PROOF OF THEOREM 1.3. We proceed by induction on $|G|$. By Lemma 2.1(a), Lemma 2.4 and Lemma 2.5, the hypotheses are inherited by $G/O_{p'}(G)$. If $O_{p'}(G) > 1$, by induction, $G/O_{p'}(G)$ is p -supersolvable, and thus G is p -supersolvable. So we can assume $O_{p'}(G) = 1$. Let $U = P \cap L$. If $|U| \leq p^e$, by Theorem 1.2, G is p -supersolvable. Assume that $|U| \geq p^{e+1}$. For any subgroup $P_1 \leq U$ with $|P_1| = p^e$, P_1 is S -semipermutable in G . By Lemma 2.1(c), P_1 is S -semipermutable in L . By

Lemma 2.8, L is p -supersolvable, and thus L is p -solvable with p -length 1. Since $O_{p'}(G) = 1$, it follows that U is the normal Sylow p -subgroup of L , and thus $U \trianglelefteq G$. Note that for all subgroups $P_1 \leq U$ with $|P_1| = p^e$, P_1 is S -semipermutable in G . By Lemma 2.1(b), it follows that P_1 is normalized by $O^p(G)$. By Lemma 2.7, U is centralized by $O^p(G_p^*)$. Let $\widehat{P} = P \cap O^p(G_p^*)$. Note that $\widehat{P} = U \cap O^p(G_p^*)$, and thus $\widehat{P} \leq Z(O^p(G_p^*))$. By Burnside's Theorem (see Theorem 5.13 of [4]), $O^p(G_p^*)$ is p -nilpotent, i.e., G_p^* is p -nilpotent. Hence G is p -supersolvable. \square

PROOF OF THEOREM 1.4. By Lemma 2.1(a), Lemma 2.4 and Lemma 2.5, it is no loss to assume that $O_{p'}(G) = 1$. Assume G is not p -supersolvable; we work to obtain a contradiction. Since G is not p -supersolvable, it follows that $O^p(G_p^*)$ is not p -nilpotent. Let $\widehat{P} = P \cap O^p(G_p^*)$. Then $\widehat{P} > 1$. Since $O^p(G_p^*) \leq L$, it follows that for any $P_1 \in \Omega(\widehat{P})$, P_1 is S -semipermutable in G . By Lemma 2.6, P_1^G is solvable. Let $M = \prod_{P_1 \in \Omega(\widehat{P})} P_1^G$. Then M is a solvable normal subgroup of G

and $M \leq O^p(G_p^*)$. Since $\widehat{P} > 1$ and for any $P_1 \in \Omega(\widehat{P})$, $P_1 \leq M$, it follows that $M > 1$. Since $O_{p'}(G) = 1$ and $M \trianglelefteq G$, we have $O_{p'}(M) = 1$. Since $M > 1$ is solvable and $O_{p'}(M) = 1$, we have $O_p(M) > 1$. Note that $O_p(M) \leq P \cap L$. Hence for any $P_2 \in \Omega(O_p(M))$, P_2 is S -semipermutable in G . By Lemma 2.1(b), P_2 is normalized by $O^p(G)$. By Lemma 2.9, $O_p(M)$ is centralized by $O^p(G_p^*)$. Recall that $M \leq O^p(G_p^*)$. Hence $O_p(M) \leq Z(M)$. Since M is solvable and $O_{p'}(M) = 1$, by Hall-Higman's Lemma (see Theorem 3.21 of [4]), it follows that $M = O_p(M)$. Hence M is centralized by $O^p(G_p^*)$. Recall that for any $P_1 \in \Omega(\widehat{P})$, $P_1 \leq M$. Hence for any $P_1 \in \Omega(\widehat{P})$, P_1 is centralized by $O^p(G_p^*)$. By Satz IV.5.5 of [3], $O^p(G_p^*)$ is p -nilpotent. This is the desired contradiction. Hence G is p -supersolvable. \square

4. Final Remarks

Theorem 1.3 and Theorem 1.4 have the following corollaries.

COROLLARY 4.1. *Let p be a prime dividing the order of a finite group G , $e \geq 2$ be an integer and $P \in \text{Syl}_p(G)$ with $|P| \geq p^{e+1}$. Suppose that $P_1 \cap O^p(G)$ is S -semipermutable in G for all subgroups $P_1 \leq P$ with $|P_1| = p^e$. Then G is p -supersolvable.*

COROLLARY 4.2. *Let p be a prime dividing the order of a finite group G and $P \in \text{Syl}_p(G)$. Suppose that $P_1 \cap O^p(G)$ is S -semipermutable in G for all subgroups $P_1 \in \Omega(P)$. Then G is p -supersolvable.*

Note that Corollary 4.1 and Corollary 4.2 generalize Theorem 3.5 of [7].

REMARK 4.3. For any odd prime p and any positive integer e , there exists a finite group G with p an odd prime divisor of $|G|$, $P \in \text{Syl}_p(G)$ and

$|P| \geq p^{e+1}$ such that for every subgroup P_1 of P with order p^e , $P_1 \cap O^p(G)$ is S -semipermutable in G , but P has a subgroup P_3 of order p^e such that P_3 is not weakly S -semipermutable in G . Hence our Corollary 4.1 and Corollary 4.2 are stronger than Theorem 3.5 of [7]. See the following example.

EXAMPLE 4.4. Let n be an integer with $n > e$, p be an odd prime and $T = \langle a, b \mid a^{p^n} = b^2 = 1, b^{-1}ab = a^{-1} \rangle \cong D_{2p^n}$. There exists $c \in \text{Aut}(T)$ such that $a^c = a$ and $b^c = ba$. Consider $G = T \rtimes \langle c \rangle \cong \langle a, b, c \mid a^{p^n} = b^2 = c^{p^n} = 1, b^{-1}ab = a^{-1}, c^{-1}ac = a, c^{-1}bc = ba \rangle$.

Let $P = \langle a \rangle \times \langle c \rangle$. Then P is the normal Sylow p -subgroup of G with order p^{2n} . Then $b \in O^p(G)$, so $ba = b^c \in O^p(G)$. Hence $a \in O^p(G)$, so $O^p(G) = T$. Hence for any subgroup P_1 of P with order p^e , $P_1 \cap O^p(G)$ is normal in G , and thus S -semipermutable in G . Let $\tilde{c} = c^{p^{n-e}}$. Consider $\langle \tilde{c} \rangle$. Note that $|\langle \tilde{c} \rangle| = p^e$. Then $\langle \tilde{c} \rangle$ is not weakly S -semipermutable in G . To see this, assume that $\langle \tilde{c} \rangle$ is weakly S -semipermutable in G , since $\langle \tilde{c} \rangle \leq \Phi(\langle c \rangle)$, it follows that $\langle \tilde{c} \rangle$ is S -semipermutable in G . Hence $\langle \tilde{c} \rangle$ normalizes $\langle b \rangle$. This is a contradiction since $\tilde{c}^{-1}b\tilde{c} = ba^{p^{n-e}}$. Hence $\langle \tilde{c} \rangle$ is not weakly S -semipermutable in G .

REFERENCES

- [1] Y. Berkovich – I. M. Isaacs, *p-Supersolvability and actions on p-groups stabilizing certain subgroups*, J. Algebra, **414** (2014), pp. 82-94.
- [2] Z. M. Chen, *On a theorem of Srinivasan*, J. Southwest Normal Univ. Nat. Sci., **12**(1) (1987), pp. 1-4.
- [3] B. Huppert, *Endliche Gruppen*, Vol. I, Springer-Verlag, Berlin, 1967.
- [4] I. M. Isaacs, *Finite Group Theory*, Graduate Studies in Mathematics, **92**, American Mathematical Society, Providence, 2008.
- [5] I. M. Isaacs, *Semipermutable π -subgroups*, Arch. Math. (Basel), **102**(1) (2014), pp. 1-6.
- [6] O. H. Kegel, *Sylow-Gruppen and Subnormalteiler endlicher Gruppen*, Math. Z., **78** (1962), pp. 205-221.
- [7] Y. Li – S. Qiao – N. Su – Y. Wang, *On weakly s-semipermutable subgroups of finite groups*, J. Algebra, **371** (2012), pp. 250-261.
- [8] N. Tang – X. Li, *On weakly s-semipermutable subgroups of finite groups*, Algebra Colloq., **21** (2014), pp. 541-550.
- [9] H. Yu, *Some sufficient and necessary conditions for p-supersolvability and p-nilpotence of a finite group*, J. Algebra Appl., **16**(1) (2017), 1750052 (9 pages), published online: April 04, 2016, DOI: 10.1142/S0219498817500529.

Received April 07, 2016; revised May 30, 2016